


CURIE'S
DOMESTIC
HOMŒOPATHY.

4s.

6 J. 3

50.70.5



The Gift of
Caroline L. Sawyer,
of Cambridge.
2 Sept., 1861.

Dr. Sam Sawyer

DEPOSITED IN
BOSTON MEDICAL LIBRARY,
BY
HARVARD COLLEGE,
LIBRARY.

Digitized by the Internet Archive
in 2010 with funding from
Open Knowledge Commons and Harvard Medical School

DOMESTIC HOMŒOPATHY.

It has been thought right to apprise the public, that pocket-cases, comprising all the medicines enumerated in this little volume, and varying in price from 40s. to 50s., may be purchased of MR. W. HEADLAND, 22, *Mount Pleasant, Gray's Inn Lane*. This enterprising Chemist has for years devoted himself to the preparation of homœopathic medicines, with a success which renders it unnecessary to have recourse to the continent for a supply of the respective remedial agents.

2

DOMESTIC HOMŒOPATHY.

5

By P. F. CURIE, M.D.

FORMERLY SURGEON IN THE MILITARY HOSPITAL OF PARIS;
MEMBER OF THE PARISIEN HOMŒOPATHIC AND GALRICAN SOCIETIES;
PHYSICIAN TO THE LONDON HOMŒOPATHIC DISPENSARY;
AUTHOR OF "THE PRINCIPLES OF HOMŒOPATHY,"
"THE PRACTICE OF HOMŒOPATHY, &c."

LONDON:

THOMAS HURST, 5, ST. PAUL'S CHURCHYARD.

1839.



C. Adlard, Printer, Bartholomew Close.

INTRODUCTION.

It is a received axiom, that the practice of medicine should rest with a class of men devoted to the arduous investigation of a science so difficult and important.

Among the non-medical public it is at all times a *desideratum* to diffuse wholesome views connected with the profession. By this means we are enabled to counteract prevailing errors, together with the innumerable prejudices which exist in every quarter. At the same time it places those who may be remote from medical aid in a position to administer means which may be suited to the emergency of the moment; and this too without participating in those fearful errors which result

from an ignorance of the acting properties of medicine.

The aim of this little volume is to place the public in possession of enlightened hygeinic rules, applicable to the various periods of life, and referrible as well to a state of health as to that of suffering. From the limited number of homœopathic practitioners, such a book is especially important in the present state of the science.

We shall point out the earliest attentions to be enforced in cases of severe acute disease, whilst awaiting the aid of the practitioner, and shall rapidly, but as lucidly as possible, indicate the treatment of acute affections which may be less serious and of more frequent occurrence.

This will enable parties, who may be resident beyond the ready access of medical assistance, either to treat themselves, or confidently submit to the direction of some intelligent friend, who may be otherwise unconnected with the profession of medicine.

This is not the place in which to insist upon the value of sound hygeinic regulations. Every individual is fully impressed with their importance, and the chapter devoted to the subject will

abundantly prove that the homœopathic hygeine is based upon firm and healthful principles.

The attentions to be observed previously to the arrival of a practitioner, and applicable to patients labouring under acute symptoms, are similarly important. In embracing the allopathic routine of medicine, we are in danger either of counteracting the efforts of nature, or of inducing vast disturbance in the animal economy. The consequences of such interference are too frequently irreparable.

With reference to the less important ailments, there is no mother of a family who can fail in appreciating the worth of a wholesome treatise respecting them; for scarcely a day passes in which she is not called upon to remedy a derangement in one or other of her children: and how is she to act without an intelligent guide? Habit will doubtlessly impel her to tread in the accustomed track: she will have recourse to purgatives; and by this means will, probably for a series of years, entail suffering upon the children she is called upon to nurture and rear.

In the following pages, meriting the attention of every mother desirous of improvement, and solicitous that her children should profit by the

advances effected in science, will be found the means of preserving and restoring their health, without subjecting them to unnecessary suffering, or to a discipline violent in its character, and frequently entailing consequences more formidable than the disease itself.

The dawn of reason, however, is at hand; and the period is not remote, when every mother will appreciate and revere the invaluable science which has been given to the world by Hahnemann.

30, BROOK STREET,

GROSVENOR SQUARE.

CONTENTS.

HYGEINE.

	Page
General Remarks	1
Attentions to be paid to Children	3
On Regimen	8
Aliments allowed—Vegetables	9
Aliments derived from the Animal Kingdom	10
Liquids	11
Aliments prohibited	12
Animal Food	13
Drinks	14
Air and Bodily Exercise	ib.
Habits	16
Observances to be enjoined on Persons suffering from Acute Diseases	17
Administration and Repetition of Medicines	19
Of Preservatives	22

DISEASES.

	Page
Aphthæ or Thrush	25
Asthma of Millar	29
Miscarriage	31
Wounds	34
Burns	36
Cardialgia	37
Head-ach	39
Colic	44
Congestion of Blood in the Abdomen	52
Chest	53
Head	55
Constipation	57
Contusions	62
Cramps—Convulsions	63
Cramps of the Stomach	66
Cries of Young Infants	71
Milk-crusts	72
Itchings	73
Dentition	76
Passion	79
Derangement of the Stomach	80
Diarrhœa	85
Pains following Childbirth	89
Poisoning	90
Swelling of the Cheeks	93
Enlargement of the Glands of the Neck and of the Ears	95
Chilblains	ib.


CONTENTS.

xi

Page

Hoarseness	96
Erysipelas	99
Quinsy	101
Croup	103
Excoriation of the Nipples	105
Children	106
Catarrhal Fever	108
Fatigue	111
Milk-fever	112
Flatulence	114
Sprains	115
Fright	116
Boils	118
Swelling of the Glands	119
Hemorrhage	ib.
Hemorrhoids	124
Catarrhal Inflammation of the Throat	127
Cerebral Inflammation	129
Inflammation of the Eyes	132
Catarrhal Ophthalmia	134
Inflammation of the Lungs	135
Breast in Nurses	138
Vigilance	140
Irritability of the Skin	141
Intoxication	142
Jaundice	ib.
Fainting Fits	143
Fetid Breath	146
Pains in the Loins	147

	Page
Acute Inflammation of the Lumbar and Psoas	
Muscles (Lumbago Inflammatoria) . . .	149
Cholera	151
Measles	155
Scarlatina	161
Small-pox	165
Megrim	169
Odontalgia	174
Otalgia, or Ear-ach	178
Palpitations	180
Whitlow (Paronchia)	181
Chicken-pox	182
Rheumatism	183
Cold in the Head	188
Bleeding from the Nose (Epistaxis) . . .	190
Stomacace (Ulcerative Inflammation of the Mouth)	192
Cough	194
Swelling of the Parotid and Submaxillary Glands	197
Worms	198
Vomiting	201



DOMESTIC HOMŒOPATHY.

OF HYGEINE.

HYGEINE is a division of the healing art which prescribes the means of preserving health. This branch of medical science will become more and more important, and will still further extend its happy influence, in proportion as intelligence and morality shall pervade the various classes of society. Not only will the judicious observance of ordinary regimen, of pursuit, of exercise, and of cleanliness, contribute to the preservation of the general health, and render epidemic diseases less frequent and intense, but still more will hygeine, as a preventive, neutralise atmospheric influence, and obviate the mischiefs resulting from humidity, or from any external causes producing

disease. Again, it must perceptibly conduce to the extirpation of hereditary taints—those poisons which, by reason of any exciting cause, such as a cold, or excess of any kind, give birth to chronic affections, multiform in character, and lamentably destructive to human life.

Homœopathy will effect rapid strides towards hygiene, since by its aid we learn that the greater portion of humanity is more or less deeply impregnated with one or other of the three grand miasmata, productive of chronic disease. These miasmata are Psora, Syphilis, and Sycosis. They are handed down from generation to generation; or they may, under certain circumstances, be communicated by one individual to another.

Conversant, in this manner, with the elementary cause of disease, and possessing a law which promises a specific for every form of malady, we may indulge a hope, in succeeding generations, to exonerate the human species from the baleful effects resulting from hereditary virus.

Children, deriving their existence from parents affected with chronic ailments, bear with them the germ of disease, which will develop itself sooner or later beneath the influence of exciting causes, such as irregularities of regimen, atmospheric changes, moral delinquency, &c.

The hygeinic attentions to children, inheriting such predisposition, should be directed with a view to combat this virus, the first grand cause of disease.

Upon this important question, we need say but little in a volume which is altogether elementary. We shall endeavour to prove, however, that the medical man ought not to be considered as competent merely to check a formidable disease, but that his counsels should serve as a guide in the *physical* education of children, and, by their hygeinic and medical attentions, should give them an unrestrained and full development of healthful life.

Attentions to be paid to Children.

From the period of birth, man cannot be regarded with too much anxiety: nothing should be omitted which may favour his physical development, prepare for him a career devoid of infirmity, and enable him to become serviceable to his family, and to his fellow-creatures, and thus endow him with one of the primary essentials to happiness.

It is remarked, however, that man is from his cradle the butt of ignorance and of prejudice, and the victim of the most baneful practices. Scarcely has the infant been launched into life, when too frequently he receives impressions which destroy or

disorder his economy for a long series of years. Recourse is had to various descriptions of purgatives, according to the climate and the custom of the nurses, to facilitate the discharge of the meconium. This practice is particularly obnoxious, from the delicate organization of the infant: it is mischievous to attempt to aid, or rather to force nature, particularly where she requires no assistance.

The secretions comprised within the intestinal canal, have at this period nothing abnormal in their character, and will discharge themselves without the aid of purgatives. Medicines should be reserved in case of need.

The temperature to which the newly-born infant is introduced, should approach as nearly as possible to that which it has just quitted: to a frame so delicate, every excess of heat or cold must be prejudicial.

The movements communicated to the child should at all times be very guarded. The practice, therefore, of forcibly rocking him is highly objectionable, since by this process he is not unfrequently rendered giddy, and exposed to cerebral congestion.

In case of vigilance, have recourse neither to the cradle nor to the exhibition of narcotics; by such practices the infant will become disordered, pine, and require medical aid.

Every circumstance should be avoided which may too sensibly affect the child; sound and light must be regulated with great caution; and it is by degrees only that exposure may be made to the noise and full glare of day.

We will offer a few hints with reference to the diet of early infancy. Most persons are conversant with what is most suitable; the milk of the mother, or, in default, that of a healthy nurse. If the latter cannot be obtained, the milk of a cow, sheep, ass, or goat, diluted with water, and rendered lukewarm, may be substituted. The milk should be prepared each separate time it may be desirable to feed the infant, and the periods should be regulated with every possible care.

During the first three or four months, the milk of the mother, or that of the cow, will in most instances suffice. Up to this period we should offer no additional nutriment. White bread, stale and grated, or biscuit-powder, boiled in milk, is now more appropriate. It is at a subsequent period only that pap, made from flower or oat-meal, the fecula of the potato, meal, &c., and prepared with milk, is admissible.

When the earlier teeth have protruded, we may in some measure vary the dietary, but it should be exclusively of a vegetable character, alternated with the preparations adverted to above.

In the earlier stage we must studiously avoid everything of a greasy nature. Milk and water should be the only beverage.

Sweetmeats and *bon-bons* are at all times objectionable, and should consequently be given with very great caution.

Animal food should not enter into the dietary until after the accession of the canine teeth, and even until the period of puberty it should constitute but a small proportion of the regimen. Lean meat is alone suitable to the digestive powers of the infant; pork, duck, &c. being wholly inadmissible.

Children digest readily, but excess of food is prejudicial: their repasts should be at short intervals, and consist rather of what is nutritious than of what is calculated to excite the appetite.

The principal repast of children should be taken about mid-day: they should not eat at too late a period, nor in too large a quantity, lest their rest be disturbed by indigestion.

Strict observance of cleanliness will contribute much to the improvement of the child: ablutions and frequent change of linen are essential; a bath, lukewarm, for five or ten minutes every day, will conduce in a striking manner to health and full development, and may be continued with advantage until the second or third year.

At a later period we may have recourse to river-bathing, during the summer season: a repetition several times in the week will prove beneficial.

Air and exercise are essential to the health and full development of infancy.

As regards attire, it should be sufficiently loose to prevent compression of the body, or restraint of motion; and we should take particular care that it be neither too heating nor too scanty.

With young girls the practice of wearing corsets is very objectionable, and should be prohibited. They are generally fashioned so as to protrude the bosom after its development, and to retain it constantly in a forced position, and in one at variance with nature. It is an error to suppose them of utility in supporting the mammæ, and in preserving the upper portion of the body in a vertical position. Instead of attaining these objects, they prevent the natural formation of the breasts, which become flaccid by their adoption. In like manner they destroy the power of the muscles of the back, and induce a species of atrophy, which renders the wearer incapable of supporting the body in an erect position. Hence result those malformations so frequently observable in young females.

The neglect of appropriate gymnastic exercise, especially with a sickly constitution, contributes

much to induce malformations; and it is no less true that the adoption of corsets, more particularly as favoring existing predispositions, is intimately connected with spinal curvature.

On Regimen.

The object of the regimen to which we subject an invalid is to dispose his system to the favorable reception of salutary medicaments. To attain this end, we must keep in observance the two following rules:

1. To prohibit the use of any substance possessed of medicinal properties, and which might thereby modify or interfere with the action of the medicines.

2. To regulate the quantity of nourishment, so that the daily loss may be supplied without surcharging the digestive organs.

These two rules form the basis upon which all hygeinic prescriptions should be founded, and they should serve equally as a guide to the regimen of persons in good health. We do not positively maintain that persons in health should avoid everything proscribed to the sick: we merely wish to observe that every substance which may produce pathological effects becomes noxious in course

of time, and should be adopted with caution in a judiciously chosen regimen, and one calculated to preserve the mind and body in a state of vigour and serenity.

The basis of the prescribed dietary we shall examine according to this principle. It will indicate the various aliments derived from the vegetable and animal kingdoms, with those which may be admissible for the patient, and those which must be prohibited during the homœopathic treatment.

Aliments allowed—Vegetables.

Man's principal nourishment is evidently derived from the vegetable kingdom, the family of graniferous plants alone supplying a large portion of his aliment. Wheaten bread is the most nutritious and the most wholesome of our food, and readily assimilates with our system.

The potato, as an article of food, is equally salutary, and the simple manner in which it is prepared is usually the best, taking care to deprive it as much as possible of its moisture.

Macaroni, vermicelli, semolino, sago, salep, arrow-root, the fecula of the potato, rice, French beans, peas, lentils, &c. as articles of food, and

prepared in a suitable manner, furnish healthful nutrition, and frequently of themselves are serviceable in convalescence from serious maladies, where the invalid should be gradually introduced to a more substantial dietary.

The melon, fully ripe, and in small quantities, may be permitted where the digestion is good.

Strawberries, gooseberries, raspberries, perfectly ripe, and prepared with sugar, or in the form of jelly—apricots, peaches, grapes, pears, and other fruits, not acid, and well ripened, may frequently enter into a well-regulated dietary.

Aliment derived from the Animal Kingdom.

Beef is one of the meats best suited to our digestive powers, and should form the basis of our animal nutriment.

Mutton may be ranked in nearly the same class as beef.

Ham, from its mode of preparation, is generally more wholesome than pork ; its adoption, however, during homœopathic treatment, must be very guarded.

The flesh of the *hare*, the *roe-buck*, and *game* in general, is in most cases very proper.

Poultry, such as pullets, capons, turkeys, &c.,

is admissible; but, unless thoroughly dressed, the flesh is too mucilaginous, and must be taken sparingly.

Grouse, pheasant, woodcock, wild duck, &c. are far more wholesome than the poultry from the farm-yard, provided they are not kept too long after being killed.

Salt and fresh-water fish are generally wholesome, and of easy digestion.

New-laid eggs, in their various modes of dressing, may be recommended with advantage.

Liquids.

There is no beverage so suitable, during homœopathic treatment, as pure water; nor is any beverage so conducive to the digestion of those who live abstemiously. In acute diseases we accordingly prescribe pure water, or *eau sucrée*, or toast-water, milk-and-water, barley-water, water-gruel, or gum-arabic water.

In chronic diseases, similar diluents are appropriate. Individuals, however, who have accustomed themselves to malt-liquor, may continue its use in moderation, provided it cause no inconvenience and be of perfectly good quality. They, too, who for a series of years have been in the

habit of drinking wine, might occasionally suffer much from its total interdiction: although such instances are rare, it is advisable to guard against them: in these cases we may permit wine in moderate quantity and diluted with water.

Tea is, in most cases, particularly objectionable as a beverage during homœopathic treatment, inasmuch as it possesses medicinal properties by no means equivocal. We should therefore for this reason be very cautious in its use. Even by those most accustomed to it, it should be taken weak, and in moderation.

Similar objections are referrible to coffee, which is even more pernicious than tea; it should therefore be still more rigorously abstained from.

Cocoa, properly prepared, is of great importance as an article of food, and should in a great measure supersede the use of tea and coffee during medical treatment.

Chocolate, taken occasionally, provided it be free from spice, is admissible as an article of regimen.

Aliments prohibited.

Asparagus, sorrel, radishes, celery, &c. should, during treatment, be particularly guarded against, as they contain properties essentially medicinal.

Cooking, in some degree, corrects the medical properties of the onion, garlic, shallot, parsley, and chervil; we should, nevertheless, exclude them as much as possible from the dietary of an invalid.

The mushroom, as difficult of digestion, should be avoided, more particularly by patients whose digestive organs may be impaired.

All fruits which are not perfectly ripe, or such as are acid, should be prohibited, since they are either difficult of digestion, or may counteract the properties of the medicine.

Chamomile, the elder-flower, and all ptisans should be prohibited.

Seasonings are referrible chiefly to aromatic substances, and are classed among the medicaments: they should therefore be excluded as much as possible from the regimen of the sick. Salt, butter, cream, and sugar, are generally sufficient to render food agreeable to the palate; it is habit alone which has created a desire for highly-seasoned dishes.

Animal Food.

Veal, as an aliment difficult of digestion, should be carefully guarded against by persons whose

stomach is delicate, and whose organs of digestion are weak.

Pork should be abstained from as much as possible during homœopathic treatment.

Ducks and geese are indigestible, and must therefore be prohibited.

If we do not altogether exclude white and mucilaginous meats, such as lamb, chicken, &c., we must at least insist upon their very moderate use.

Drinks.

All drinks too exciting in their character, or possessing medicinal properties, should be expunged from the homœopathic regime. These stimulants comprise strong wines, spirits, tea, coffee, &c.

The regimen should at all times be appropriate to the age and temperament of the patient; and from the rules already laid down, we may easily determine upon the regimen adapted to each individual.

Air and Bodily Exercise.

It will be superfluous to dwell upon the importance of pure air, all persons being fully alive to its beneficial effects upon those who are ill: the pa-

tient should enjoy it upon all possible occasions. We are fully aware, too, how essential is bodily exercise to the regulation of our natural functions. It preserves the equilibrium of the system, and corrects, in a striking degree, the mischief produced by a too liberal table, the result of our social habits. Most persons eat and drink more than suffices for the support of life; it is fortunate, therefore, that exercise has the power of correcting the inconveniences arising from improprieties of regimen.

In chronic affections, exercise in the open air is indispensable; it induces an important and salutary action in the muscular system, and obviates visceral congestion.

Walking in the open air is most beneficial for the invalid; he should regulate his walks according to his ability, and never prolong them so as to induce exhaustion.

Since walking and dancing generally call into action the lower extremities only, it is important, especially for the young, to have recourse to those gymnastics in which the upper extremities may participate.

Riding on horseback, or in an open carriage, is advisable where the patient may be incapable of walking; but the latter exercise, in connexion with gymnastics, is greatly to be preferred.

Habits.

Habits, such as late hours, the attendance at balls, concerts, theatres, the gaming-table, &c., from their exercising an important influence upon the successful or unsuccessful issue of disease, must not be passed over in silence. Chronic cases, by such indulgence are frequently rendered incurable. The patient stands in need of regular repose and serenity of mind, to obviate any interference with the favorable reaction to be brought about by medicine. He must sedulously avoid all the indulgences we have just denounced.

It is important that the invalid should accustom himself to light clothing, lest an undue warmth should render him susceptible of colds, excite perspiration, and debilitate the system. The invalid, however, should be careful in protecting himself against the influence of cold, and more especially against changes of temperature: our remarks refer merely to excess in the use or in the neglect of clothing.

He should avoid perfumes, both on the person and in his apartments, and should abjure an excess of sweet-smelling flowers: they exercise a deleterious influence upon homœopathic remedies.

The use of cosmetics is similarly objectionable. Tobacco should be prohibited, unless the patient have acquired a confirmed habit of taking it, and where the sudden and total interruption of its use might be prejudicial to his health. In all habits of this class, such as drinking of wine, tea, coffee, &c., we must gradually diminish the quantity, and not abstain entirely, and on the instant.

The indiscriminate use of warm baths is pernicious, and must be strictly prohibited; cold or tepid ablutions are preferable during medical treatment.

*Observances to be enjoined on Persons suffering from
Acute Diseases.*

The hygeinic observances in acute affections are rest, bodily and mental tranquillity, mild and equable temperature, and the avoidance, so far as practicable, of such causes as may have given origin to the disease.

It is generally observable, in acute cases, that the greater part of the functional economy is deranged. In most instances the appetite is impaired, or entirely suspended. The failure commences with aversion for food; and nature thus, of

her own accord, indicates the observance of the first hygeinic rule. To an invalid entirely devoid of appetite, no nourishment whatever should be offered; instinct should serve as a guide, and be implicitly followed. In adopting this plan, we need be under no apprehension of debilitating the patient, for the malady is brief in duration, and the functions are speedily restored. The appetite and digestive powers resume their natural course, and the affected organs are quickly re-established. More particularly we must bear in mind that the invalid take nothing which may be considered strengthening; he should receive only what he may have the power of digesting. In cases where the digestive functions do not act, or act imperfectly, the superfluous food stimulates, to a painful degree, an organ which should remain quiescent; and, far from strengthening the patient, throws him into a state of great debility.

Regimen, or positive abstinence, is not in all cases, or for a long period, essential in homœopathic treatment; in the more serious ailments, we may prescribe, after a brief interval, light nourishing food, gradually proportioned to the restoration of the economy. In a very short time the patient may have recourse to his ordinary regimen.

In less important affections, where the appetite

is not suspended, we should carefully regulate the diet, and recommend the moderate use of such aliments as may be suitable.

As regards the medical attentions applicable to acute diseases, we shall point them out in the chapters dedicated to the respective maladies. We shall dwell upon the more frequent and less serious affections, giving a glance merely at those of greater magnitude, and which should rest as much as possible in the hands of the physician.

Administration and Repetition of Medicines.

The homœopathic remedies are administered in globules, in tincture, by olfaction, and externally by means of friction.

The globules are placed upon the tongue, or, being bruised, are taken in a little sugar of milk; or again, they may be dissolved (each) in one or more spoonfuls of pure water.

If it be advisable to administer only a portion of a globule, it may be dissolved in as many spoonfuls of water as may correspond with the required divisions of the globule; and, pursuing this plan, we may exhibit several spoonfuls, or fractions of a globule, in the course of the day.

The tinctures are exhibited in solution with

water. The prescription of an entire drop, in the course of the day, is seldom required; consequently, when we have recourse to tinctures, their administration is virtually the same as that of the globules dissolved in water.

When olfaction is advisable, one or two globules of the selected remedy should be placed in a small phial, and its aperture be applied to the nose of the patient either for a few seconds or for a minute or two, as the case may require.

The use of medicaments externally, whether as lotions or as frictions, has been hitherto but rarely practised. Experience, however, will doubtlessly point out many cases where this mode may be preferable. At present we derive much valuable assistance from arnica, as an external application in contusion, from heated oil of turpentine, or from alcohol, in cases of burns.

In most instances the medicine should be administered in very minute doses, one, two, or three globules, or even the fraction of a globule, according to the susceptibility of the patient.

With reference to the most eligible dilution, the highest should in most instances be selected; nevertheless, as regards this great practical means, it is the choice of a suitable remedy which chiefly avails. A judicious election once made, we may succeed perfectly well with dilution, 12, 18, 24,

or 30, generally selecting the higher number for the more susceptible patients.

As the more suitable method of exhibiting the remedy, it appears to me that the globules or tinctures, diluted with water, are preferable; but it is evident that perfect cures have been effected by other modes.

These signs will assist us in designating the dilution of the doses: $\frac{0}{30}$ signifies a globule of the 30th attenuation; $\frac{1}{x}$ has a similar signification. The superior figure or sign points out the number of globules; and the figure, whether Roman or Arabic, placed at the right or inferior part, indicates the degree of attenuation.

6 corresponds to	II. or a billionth.
9	III. or a trillionth.
12	IV. or a quadrillionth.
18	VI. or a sextillionth.
24	VIII. or an octillionth.
30	X. or a decillionth.

As regards the repetition of the dose, it is evident that each case, and each separate constitution, must present variations as numerous as the circumstances indicating the dose itself. In slight affections, however, one, two, or three doses will generally suffice; and we find that one or two doses in the course of the day, will, in

most instances, meet the case. In the more serious cases, we must have recourse to practitioners of acknowledged repute. Upon this important subject, which is not within the province of an elementary volume, we must refer the reader to the author's "Practice of Homœopathy."

It is to be remarked, that in all instances where improvement is visible, we should never interfere with the favorable reaction; on the other hand, where the medicine produces no effect, it may be repeated; and where it has been injudiciously selected, we must substitute one more appropriate.

In cases where it is evident that the medicine acts too powerfully, and causes suffering to the patient, we must exercise due caution in repeating the dose. We must carefully note the result, and if the patient suffer much, we should administer a suitable antidote. As regards the exhibition of antidotes, we are guided by precisely the same rules as in the administration of other remedies.

Of Preservatives.

No one, in the present day, can for a moment doubt the importance of Vaccination as a preservative against Small-pox. Notwithstanding

the early prejudice against its introduction, it has gained the full confidence of an enlightened public; and the value of the discovery is every day more fully appreciated.

The preventive agency of the vaccine virus enters perfectly into the spirit of homœopathic medicine. The virus acts as a preventive, by causing an eruption similar to that of small-pox. The advantage we have gained in variola, we may yet hope to obtain in other diseases; and homœopathy has already put us in possession of certain preservatives, which I shall immediately proceed to point out.

A child may be protected against SCARLET FEVER, the *Scarlatina Mitis* of Sydenham, by causing him to take *Belladonna* during the period of the epidemic. One or two globules may be administered every four or five days, and from five to eight doses will usually suffice for the protection of the child. In cases where the disease is already apparent, the exhibition of belladonna will render it mild and trivial in effect.

Against the MEASLES, *Pulsatilla*, alternated with *Aconite*, will act as a preventive. We commence with the administration of one or two globules of *pulsatilla*, and, on the following day, we may give one or two globules of *aconite*: this treatment should be repeated three or four

times. The same medicaments will serve to modify the disease, if it have already presented itself.

IN MILIARY FEVER we may exhibit *Belladonna* and *Aconite*, alternately, as a preventive, during the period of its prevalence. We begin the treatment by giving one or two globules of aconite, and after a couple of days we give belladonna in a similar dose, repeating the medicines three or four times.

Camphor has been administered as a successful preservative against CHOLERA—one drop of the tincture being taken each time on approaching the sick, or upon entering an infected place. The administration should be alternated with *Veratrum* and *Cuprum*, in the proportion of one or two globules, every four or five days.

OF DISEASES.

Aphthæ or Thrush.

APHTHÆ are vesicles situated upon the mucous membrane which lines the cavity of the mouth. These vesicles are either isolated, or confluent to such an extent as frequently to occupy a great portion of the mouth.

Precursory Symptoms. Great uneasiness, vigilance, difficulty of breathing, and of taking the breast; breath fetid; the voice hoarse and feeble; tongue dry and swollen; dryness, heat, and redness of the mouth and throat; exhaustion and somnolency; dulness of sense.

Symptoms. Small points or elevations, of a palish or grey colour, subsequently make their appearance; these gradually enlarge, and form superficial pustules, round, about the size of a millet seed, and filled with a sanious fluid.

The viscid mucus discharging from these pustules very rapidly forms little scabs, which fall off either immediately, or after a few days, and are very quickly renewed.

These aphthæ are of trivial moment when they are white, of a pearly grey or yellowish aspect, transparent, moist, diminutive, easily crushed, isolated, and disposed to fall off readily and quickly, and affect merely the mouth. Frequently, however, they become dangerous, and are then of larger size, dull, blue, grey, brown, livid, gangrenous, red, fetid, numerous, compact, confluent, covered with thick and very adhesive scabs, or entirely dry and hard; are frequently renewed, and readily affect the pharynx, intestinal canal, and even the windpipe. The aphthæ usually affecting infants at the breast are very rarely malignant. It is of these last mentioned that we especially treat, since the form adverted to above is in most instances symptomatic of more important affections.

The most frequent sources of aphthæ in young children are, want of cleanliness; food, prejudicial in quality, or ill suited to their age. In many instances, too, they originate in improper things which they are permitted to suck; or from the sucking-bottle not being properly cleansed after each period of use.

Treatment. *Borax* is one of the most suitable remedies in aphthæ of infants at the breast,—dissolving a globule in ten teaspoonfuls of pure water, and giving three, four, or five spoonfuls

a day, according to the result: the child usually recovers in a very few days.

Mercury is at times indispensable; it is especially indicated for children of a gross habit, and where the following symptoms are remarked: viz.

Aphthæ, after slight febrile excitement, extending themselves over different portions of the lining membrane of the mouth; where the whitish aphthæ are converted into slight ulcerations with red edges; are white or yellow, with burning pain, frequent bleeding, and difficulty in swallowing; pallor of countenance, puffiness of the cheeks; swelling of the lips, with enlargement of the submaxillary glands; salivation, fetor of the mouth, resembling the mercurial odour; whitish tongue; bowels irregular, alternately confined and relaxed; urine thick and clouded; dry cough, great uneasiness, debility, and vigilance.

One or two globules of mercury in four ounces of water,—two tablespoonfuls every day will speedily relieve this train of symptoms. Should no change be perceptible, the dose may be increased.

Sulphur will be found of great value in cases of complication, and is especially indicated where we remark a psoric tendency; where the mouth, lips, and tongue are incrustated with thick scabs; where the evacuations are green, frequent, and attended with tenesmus; where the body is covered

with a rash; where the thighs are excoriated; and where the back is affected with a vesicular eruption.

Arsenic is especially indicated where the aphthæ have assumed a malignant character.

The medicines which I have pointed out are not the only ones which are valuable, but they are the remedies which are most frequently had recourse to. Some particular circumstances, however, may indicate the propriety of other agents. Of these we shall subjoin a list, which the practitioner may consult in case of the inadequacy of the preceding, or in consequence of any special indications:

Acidum Muriaticum; Acidum Nitricum; Acidum Sulphuricum; Agaricus; Ambra; Ammoniæ Carbonas; Ammoniæ Murias; Anacardium; Argentum; Baryta; Bryonia; Calcareæ; Cantharides; Capsicum; Carbo Animalis; Causticum; Chamomilla; China; Daphne Mezereon; Dulcamara; Graphites; Helleborus; Iodium; Kali Carbonas; Kali Hydriodas; Kali Nitras; Lycopodium; Magnesia; Magnesiæ Murias; Manganese; Natrum; Natrum Muriaticum; Nux Vomica; Phellandrium; Phosphorus; Plumbum; Pulsatilla; Rhododendron; Sepia; Spigelia; Spongia; Staphysagria; Strontium; Thermæ Tœplitzens; Thuya.

Asthma of Millar.

This is an infantile disease, analogous in symptoms to those of croup: hence it is that these two diseases have frequently been confounded, both by parents and by practitioners of slender experience. This affection is sudden in its attack, and is preceded by no premonitory symptoms; it exhibits itself more particularly in children, between the ages of two and seven years, and in the winter season, after having contracted a cold. The first indication is a sense of choking, great anxiety, with difficult respiration; the voice becomes harsh, and acquires a deep tone: should there be cough, it is hoarse, broken, slight, infrequent, dry, and generally devoid of mucus. When the affection is very severe, it is accompanied by heat of skin and frequency of pulse, and perspirations supervene upon the close of the attack. Unless the first attack prove fatal, it usually terminates in the course of a few hours in sneezing, eructation, and vomiting; the child sleeps tranquilly, but on awaking is weak and languid, and his symptoms very closely resemble those of catarrh. A fresh paroxysm, more serious than its predecessor, comes on within twelve or twenty-four hours.

The most general remedy is *Sambucus*, particularly where the patient falls into a state of somnolency, with the eyes and mouth half-opened; is unable to inspire freely; is compelled to sit down; and where, even in that position, the air enters the lungs with a whistling sound, accompanied by a sense of suffocation; he strikes his hands together behind him; the hands and face become swollen and purple; there is heat of skin, without thirst; the child weeps at the access of the paroxysm; where there is cough; and where the attack comes on about the middle of the night.

We exhibit, then, one or two globules of *Sambucus*, to the little sufferer, and carefully wait the result: if no effect be apparent, the medicine may be repeated in one or two hours' time. Several doses may, in this manner, be required to arrest the progress of this formidable disease.

Moschus is serviceable in this affection, when the patient feels as though he had inhaled the vapour of sulphur.

Ignatia, *Nux Vomica*, and *Pulsatilla*, should, under particular circumstances, come under the consideration of the practitioner. In consulting the *Materia Medica Pura*, he will ascertain from the symptoms which of these medicines

should follow the exhibition of the *Sambucus* or *Moschus*, should the latter remedies fail in the removal of the disease.

Miscarriage.

By miscarriage we understand an accouchement taking place during the earlier months of pregnancy. The causes of this accident are the psoric condition of the patient; the weakness of the mother; a peculiar irritability of the womb; predisposition, resulting from prior miscarriages; mechanical lesions, such as disturbance of the lower parts of the abdomen, from a blow, a fall, or violent fit of passion, grief, vexation, anger, or other moral affection. Sometimes a miscarriage takes place suddenly, and without intimation; in other instances, it may be announced by previous indisposition. The earlier symptoms are violent pains in the uterus, drawing and cutting pains extending from the kidneys towards the interior of the pelvis; shiverings; great desire to pass water; with viscid and sanguineous discharge from the vagina. Should this discharge increase, and the pains be not readily checked, miscarriage will be almost inevitable.

Treatment. The patient must strive to main-

tain the greatest tranquillity both of mind and body, and must assume the recumbent position.

Chamomilla, of the 12th dilution, should be administered where the symptoms, as detailed above, shall present themselves.

Belladonna, at the 30th attenuation, is indicated where the pains are violent, tensive, and pressing throughout the entire lower portion of the abdomen, and more especially towards the pubic region; where there is depression of the reproductive organs, and violent pain in the region occupied by the kidneys.

Hyoscyamus is beneficial in cases where we meet with convulsions, and diffused cramp-like pain, succeeded by stiffness of the joints, and accompanied by loss of sense.

Ipecacuanha is the most appropriate remedy, where the patient is seized with tonic cramp, stiffening the entire body, and conjoined with loss of sense.

Crocus, of the 3d attenuation, is serviceable where the sanguineous discharge is black and clotted; where the pains appear to pass through from the umbilicus to the womb; and where there is pain in the vertebral column.

Sabina, administered in the early stage of pregnancy, will act as a preventive in those who may have a disposition to miscarry.

Nux Vomica is valuable in cases where the predisposition results from a varicose state of the reproductive organs, originating either in the abuse of coffee, or in obstinate constipation, which, from inordinate pressure, induces premature contraction of the uterus.

Carbo Vegetabilis and *Lycopodium* are serviceable in similar cases where the psoric tendency is more or less developed.

In cases where there may be a disposition to psora, but without a varicose tendency, *Sepia* and *Silicea* should receive the preference.

Arnica is prescribed with success where abortion is to be apprehended from a fall, or from external violence: we may use it also as a lotion to the injured or contused surface.

Asarum, Cannabis, Cantharides, China, Nux Moschata, Platinum, Plumbum, Rhatania, Calcarea, Ferrum, Secale Cornutum, Zincum, and Sulphur, may in like manner be employed in the prevention of miscarriage.

In declaring the principal remedial agents in this affection, we invite the patient to treat herself in a case of such vital import, and where she may be distant from a homœopathic practitioner. We specify the disease from the prompt assistance required for its successful treatment.

Wounds.

In serious wounds we must have recourse to a surgeon of acknowledged repute. In cases presenting but little danger, the following treatment should be adopted. After carefully washing the part, the divided edges are to be brought into apposition, and kept moistened with the tincture of arnica. When union has been effected by means of a suitable plaster, we place over the part a compress steeped in arnica, support the whole with a moderately tight bandage, and administer arnica internally.

The traumatic fever either does not declare itself, or is at least very sensibly moderated by the use of arnica, applied both externally and as an internal remedy. The dose of the tincture in water is in the proportion of from 10 to 20 drops to the ounce. We administer one, two, or three globules during the first four and twenty hours, and, should occasion require, we may have recourse to many repetitions of the medicine.

When the wound becomes very painful, either without any appreciable cause, or from one which we can remedy, such as a bandage too tight, the presence of a foreign body, &c., the patient at the same time feeling much excitement, excessive

sensibility, a disposition to tears, agitation, or vigilance, we should exhibit a dose of *Coffea*, some globules or even a part of a drop of the tincture. Should the *coffea* prove inefficient, we may follow up the treatment by a dose of *China*, which will complete the cure.

If the wound be circumscribed by an extensive red areola, and if the surface be swollen and inflamed, *Pulsatilla* will generally produce beneficial results. We administer one, two, or three globules, according to the excitability of the patient, and in most instances select it from the 12th up to the 30th attenuation.

Mercury will very speedily check any disposition which the wound may have to suppurate.

In the treatment of wounds we should scrupulously avoid the use of all medicinal ointments which are generally in so much request. To prevent any portion of the apparel attaching itself to the skin, it would be well to shave the hair, and apply small bandages spread with some greasy substance: simple cerate, or cocoa-nut oil, is very suitable.

If the wounds are very extensive, and attended with loss of substance, in addition to the bandages which cover the edges of the wound, we may apply an open compress slightly spread with cerate: we may afterwards impose a fold of lint.

In these cases we must guard against the exhibition of arnica in too concentrated a form, since it may counteract the desired effect: 5 or 6 drops to an ounce of water will be sufficient. The wound must not be exposed more than once during the day, and where there is no suppuration, there will be no need of disturbance until the cure is complete.

Burns.

Burns, as all persons are aware, present various degrees of importance, and exhibit themselves in three distinct stages. In the earliest stage the part affected is inflamed, the skin is puffed up, and blisters are formed; in the second a portion of the skin is more or less deeply destroyed; in the third stage, not only is the skin destroyed, but also the subjacent structure.

It is principally the two former stages which will be considered here; the last requiring the especial regard of an experienced practitioner.

Should the pain be limited in extent, and trivial in its nature, we may derive much benefit from allowing the parts to approach the fire. This greatly increases the pain for the first few moments, but with courage to persist, the pain

very quickly ceases, and the vesications do not present themselves. When the pain is more extensive, we may moisten the parts with heated spirits of wine, or brandy, and we must be careful to keep the compresses constantly wet. Turpentine may be used instead of alcohol or brandy; and to prevent the too rapid evaporation, the moistened cloths should be covered over with cotton. Serious burns require the unremitting attentions of a nurse, who should never allow the injured surface to become dry.

At night we should withhold the candle from the patient, or place it within a lantern, lest the alcoholic vapours should ignite.

When the burn induces febrile action, we administer one or more doses of aconite, from the 18th to the 30th attenuation.

Soap, used for domestic purposes, administered internally, quickly removes the pain, and induces a rapid cure, without having recourse to any other remedy.

Cardialgia.

The disease which has received this appellation is one having its seat in the stomach, or in the diaphragm, and frequently extending itself over a large portion of the abdominal viscera.

This affection presents itself in different degrees and forms, and necessarily requires the assistance of the medical man. In this place we shall consider the disease merely in its simplest character; the result frequently of cold, and attacking irritable and sensitive subjects, women, and children.

The region of the stomach, and frequently also the lower part of the abdomen, are swollen, sensitive, and painful; the patient experiences palpitations of the heart, anxiety, constriction across the chest, difficulty of respiration, pains in the head, a disposition to faint, frequent rising of the food, sometimes acid and always disagreeable; very rarely flatulence.

Children affected with cardialgia, are agitated and restless, have shortness of breath, are fretful, and flex the legs upon the body. It is generally induced by cold, vexation, acid wine and food, beer, fat meat, &c.

One of the most certain remedies is *Chamomilla* of the 12th attenuation. We administer one, two, or three globules, according to circumstances; and the remedy is repeated after a few hours' interval, or after twenty-four hours, if required. The physician, however, may vary the medicines according to the shades of variety in the disease, and he will generally select them from among the following:

Nux, Chamomilla, Belladonna, Cocculus, Ipecacuanha, Stannum, Staphysagria, Bryonia, Pulsatilla, Ignatia, China, Bismuthum, Arsenicum, Plumbum, Argilla, Carbo Vegetabilis, Calcareo Carbonica, Baryta Carbonica.

Headach.

Cephalalgia, or headach, is an affection very rarely idiopathic: it is generally symptomatic of a more complicated malady, which may be either acute, or classed among the chronic diseases. We shall nevertheless enumerate a few instances where the disease is sufficiently simple to render the attentions of the physician unnecessary.

Aconite is especially indicated in pains of the head, excited by determination of blood to the part, and where the following symptoms present themselves:

Aching pains in the head, vertigo, giddiness, and pricking sensation in the forehead; burning heat in the brain, whilst the body and limbs are rather cold than otherwise; pressive pain with throbbing in the head, which subside on lying down, and are aggravated on rising; the cheeks flushed and tumid, great nervous excitement,

irritability of temper, incapacity for exertion, vigilance, relief on going into the air.

Belladonna is suitable when there exist great sensibility of the scalp, and even of the hair, swelling of the veins of the forehead and of the hands, buzzing in the ears, a mist before the eyes, hypochondriacal disposition, absence of thirst.

It is also valuable where there are shiverings in the back and shoulders, in the morning, followed by a sense of uneasiness all over the body, succeeded by nausea, with inclination to vomit; pain in the head, pressure and drawing towards the forehead, and becoming insupportable upon the slightest movement of the eyes; aggravation from the slightest breath of air, and from every motion, rest alone assuaging the pain; where the pain appears concentrated in the forehead; where there is a burning spot, with pain, in the bones of the nose; aggravation from light, excessive irritability; palpitation of the heart resounding through the head; unnatural craving for food; oppression at the stomach, especially after eating.

Bryonia may be exhibited in pains of the head produced by rough weather, and by painful emotions of the mind, and when the following symptoms are apparent:

Cephalalgia continuing from morning till the evening, contraction affecting the whole of the

head, when the sufferer is compelled to lie down, and is nearly deprived of his senses; the eyes glassy, dull, and contracted, the face pale; incapability of enduring either light or noise; disposition irritable and querulous; in the afternoon, palpitation of the heart, tension across the chest, uneasiness, inclination to vomit, retchings, constipation, deafening pain in the head the next day, or even a state of fever.

Pulsatilla is applicable to pains in the head, characterized by vertigo, as from inebriety, with paleness of the face, anxiety, inclination to shed tears, palpitation of the heart, bleeding at the nose, sensibility to light, failure of appetite, &c. The same remedy is suitable when the affection of the head is semi-lateral, diminishes on walking about, and increases on sitting down.

Cephalalgia with shooting and throbbing in the left side of the forehead, in the morning and evening, relieved by pressure and the open air; aggravation in a room, on lying down, in lowering or raising the eyes; acid or bilious vomitings, after violent pain in the stomach; abdominal pains, as from gripings, colic, contraction, &c.

Cephalalgia, with shooting pain at the top of the head, burning in the eyes, inclination to fall forwards, pain in the stomach, face flushed, &c. At the lightest movement, the patient feels as

though his head would detach itself from his shoulders; melioration on compressing the head, debility after the paroxysm.

Cephalalgia, the consequence of keeping late hours, debauchery, and the abuse of wine and spirituous liquors, has for its characteristic symptoms, a pain resembling that of a fracture, heaviness, disposition to vomit, heaving of the stomach, chills, weakness of the limbs, ill humour, irritability. *Nux Vomica* is particularly suited to this train of symptoms.

In cephalalgia produced by the use of coffee, *Nux Vomica* is very efficient, particularly when the following symptoms are remarked:

Cephalalgia in the morning compelling a recumbent position, and increasing towards the middle of the day; much confusion in the head on awaking; repugnance to food; darting pain over the left eye, with pressure, and affecting the organ itself; increase of pain from the pressure; contraction of the eyelids, dread of light, dimness of vision, discharge of hot tears, heat around the eyes.

Obstruction in the left division of the nose, sweats through the violence of the pain, occasional vomitings, derangement of the reflective faculties during the pain, incapacity of thought after the pain, forgetfulness, great irritability,

reluctance to walk, noise and conversation an annoyance.

Nux Vomica is again serviceable in cephalalgia produced by constipation of the bowels.

Moschus is applicable to the cephalalgia of hysterical and hypochondriacal subjects, more particularly when the under-written symptoms are observable:

Compression above the root of the nose, with confusion of idea; vertigo upon the slightest movement of the head, compression in the head, giddiness with threatened loss of sense, pain throughout the head with twitchings at the nape of the neck, when the pain becomes tensive; determination of blood to the head, disposition to sleep, anxiety, and ill humour.

In cases where *Moschus* has failed of success, we must employ *Nux Vomica*, which will then relieve the pain.

Chamomilla is a suitable remedy when the headach results from the abuse of coffee, and having frequent recourse to *Nux Vomica*. It is of value in cephalalgia, characterized by the following symptoms:

Pain, pressive and stunning when sitting down and reflecting; weight and sense of cracking; pain even during sleep; cephalalgia in the morning, the eyes being still closed, and in a state of semi-

sleep, the pain in the head ceasing on being suddenly awakened, and when the patient has risen; pain from laughter, twitchings, &c.

Chamomilla is again admissible in cephalalgia resulting from a cold, and characterized by tearing pains in the head and shootings in the temples, great heaviness over the root of the nose, as though threatening a cold in the head; yellowness of the forehead; puffiness of the face and of the eyes; dimness of the visual organs.

Colic.

Colic, or Gripes, is a disease attended with grinding, tearing, and tensive pains, without any inflammatory symptoms, having neither fixedness nor continuity, and affecting the interior of the abdomen, but more especially the umbilical region. This pain generally directs itself from above downwards, and is frequently accompanied with swelling of the lower part of the abdomen. Sometimes we observe a kind of sinking and constriction in the region of the umbilicus, and occasionally the superior portion of the abdomen becomes swollen, more hot, and painful. The stomach is generally free from pain, and the patient

rarely vomits; if sick, he throws up his food, together with a small quantity of bile. Sometimes the complaint is attended with constipation, but more frequently with slight diarrhœa, and there is very rarely fever.

In children, after a discharge of urine more copious than ordinary, we may occasionally remark considerable agitation and impatience, continued crying, sudden screams, contortions during sleep, vigilance, and cramps; they flex the legs upon the belly, appear as though they were bent double, and are unable to suck, being either stretched out or lying down.

The treatment will vary according to the circumstances producing the disease. Among children, it frequently originates in a cold, or in worms; in adults, it may proceed from cold, excess in eating or drinking, and not unfrequently from a disposition to flatulence, and from hemorrhoids.

Chamomilla or *Pulsatilla* is the remedy most suitable in cases of colic presenting the following symptoms, and where the origin has been a cold:

Violent cutting and tearing pains in the abdomen, occasioning incessant restlessness; sensation as though the abdomen were entirely empty, with constant rumbling in the intestines, eyes appearing discoloured, accumulation of saliva in the mouth, contusive pain in the kidneys, nausea, disposition

to vomit, diarrhœa, evacuations green, watery, and mixed with mucus.

Nux Vomica is indicated where the following characteristic symptoms are observable:

Hardened fæces or constipation, sensation of a heavy weight in the lower part of the abdomen, with borborygmus; unusual heat, considerable tension of the belly, with respiration anxious, short, and laboured; fulness of the abdomen; sense of fulness beneath the ribs; pain drawing, pinching, and compressed, as though the intestines were squeezed in different directions by suffering; pressure at the pit of the stomach; diffused head-ach, with confusion of idea; the abdomen painful to the touch; loss of recollection; coldness of the extremities at the time when the affection is most violent.

Windy colic, having its seat in the inferior portion of the abdomen, yields to the same remedy; when the flatulence, in seeking to escape, produces pressure, as from a sharp or cutting instrument, to the bladder, rectum, or organs of generation; pressure, causing the patient to pause after each step, when in an erect posture, or which is quickly removed when lying down or sitting in a curved position.

Mercury and *China* are successfully employed in colics produced by worms. We generally observe the following symptoms:

Great inclination to vomit ; risings of the food ; increased secretion from the salivary glands ; torsion of the abdomen ; a sensation of hardness around the navel ; convulsion in the abdominal muscles ; pricking in the throat ; frequent hiccough ; occasional bulimia ; aversion to sweet food ; frequent desire to relieve the bowels ; the lower abdomen hard and swollen ; tensive and burning pain round the navel ; diarrhœa ; slimy evacuations ; general debility ; bad complexion ; discoloration of the eyes ; itching of the nose ; dilatation of the pupils ; itchings at the rectum.

China is chiefly applicable to children, more particularly where they pick the nose, when the eyes are dark beneath, where the sleep is disturbed, when they scream out, or exhibit any of the symptoms enumerated above.

Excess of feeding, or unwholesome articles of food, frequently give rise to colic ; we then remark the following symptoms :

Nausea ; weight ; sensation of stretching, and painful tension in the lower part of the abdomen ; severe pinching ; cutting pains in the abdomen, increased by pressure ; salivary secretion, white and frothy ; sharp tearing pains above the navel ; diarrhœa, the evacuations green or of a yellowish citron colour attended with violent pain in the stomach ; complexion pallid ; eyes dark beneath ;

slight convulsion of the limbs; curvature of the whole body; pressure and tensive pain in the head.

A cup of coffee without milk will disperse these symptoms in persons who have not accustomed themselves to its abuse. Frequently after its exhibition we shall obtain an operation from the bowels, or vomiting of the indigestible food. One, two, or three globules of *Pulsatilla*, administered at this juncture, will remove all trace of the indisposition.

The Hemorrhoidal Colics are distinguishable by the following symptoms:

Throbbing at the pit of the stomach; very painful sense of tightness and constriction, as though everything were too full; borborygmus in the lower part of the abdomen; flatulence which cannot discharge itself; distressing heat in the lower part of the abdomen, as if resulting from an effervescence of the blood; swelling of the belly; diffused heat, with swelling of the veins of the hands and forehead; the patient is unable to bear any clothing; his sufferings are greater when in bed, and amusement affords him some relief; pain in the kidneys as though they were bruised; difficulty of rising up; pain in the lower part of the abdomen, as from the receipt of a blow; agitation, anxiety, and vigilance. These symptoms are dispersed by the aid of *Pulsatilla*. We may dissolve

one, two, or three globules, according to the sensibility of the patient, in eight ounces of water, and administer two tablespoonfuls of the liquid every morning. If one dose prove insufficient, we may have recourse to a second.

Nux Vomica is often applicable in Hemorrhoidal Colics, especially when the pains proceed to the bladder, causing a continual inclination, but want of power, to pass water; where there exists spasmodic stricture of the urethra, and where the belly becomes very sensitive.

In such cases, then, we administer *Nux Vomica* in the same manner as *Pulsatilla*, taking care to exhibit it in the evening in preference, provided there be no necessity for immediate treatment.

Colocynth, of the 30th attenuation, is very efficacious in different varieties of Colic, and more particularly where the following symptoms are presented:

Violent pains in the abdomen, unremitting or returning at short intervals, and with renewed intensity. After their complete disappearance, these pains present the peculiarity of leaving, throughout the lower abdominal region, a sensation of bruising, as though the intestines were suspended by threads easily broken, and obliging the patient to walk slowly, and with extreme caution.

The suffering experienced in the superior portion of the abdomen occupies chiefly one single point, of slight compass, and returning only at intervals of five or ten minutes, or at all events very rarely. It commences most frequently in slight twitchings of the ribs, and directs itself towards the centre of the abdomen; becomes cutting, pressive, searching, tearing, and drawing, and attains such violence as to induce the patient to utter loud cries, and to bite whatever comes within his reach. He suffers much anxiety, the pain not permitting him to rest in quiet, but compelling him to twist about like a worm, and frequently causing him to be bathed in sweat.

Colics with which many women are affected at the commencement of their monthly periods, very frequently disperse when menstruation has taken its course; but when the pain continues, it is advisable to meet the case with suitable treatment.

Nux Vomica will apply where the following symptoms prevail:

A kind of twisting in the lower abdominal region, with nausea; searching pain in the same part; a sensation as from a strain; piercing, spasmodic pain in the pelvis, for the most part affecting the pubes and the groin; pain as from

a bruise ; a squeezing in the region of the bladder ; violent sense of extension in the abdomen as though it would burst.

One or two globules of *Nux*, of the 24th or 30th attenuation, in six ounces of water, and a spoonful taken every hour, will readily assuage the pain.

Coffea, of the 3d attenuation, is suitable in the following symptoms :

Pains in the abdomen, extremely violent, spasmodic in their character, and extending even to the breast ; pain as though all the intestines were divided with a knife, and which deprives the patient of her senses ; dreadful curvature of the body, with convulsion of the limbs ; grinding of the teeth ; frightful cries ; general coldness ; cramp-like sufferings ; loss of breath ; repeated sighs ; feeling as though the abdomen would burst ; fulness of the lower region of the abdomen ; pressure.

Pulsatilla may be exhibited when the menses are accompanied by violent and pressive pain in the inferior portion of the abdomen, and in the kidneys ; painful drawing sensations from the trunk to the thighs, relieved when sitting down, and which are conjoined with a painful pressure in the rectum, pains in the bones, and numbness of the thighs.

Belladonna is indicated where there exists a sensation of pressure in the genital organs, from within to without, as though these organs were striving to escape outwards.

In addition to the medicaments indicated above, as tending to check the colic, we may exhibit the following :

Alum, Asarum, Arsenicum, Calcarea, Carbo Vegetabilis, Cocculus, Hyoseyamus, Lycopodium, Phosphorus, Secale Cornutum, Sulphur, Veratrum, Conium, Capsicum, Crocus, Cuprum, Euphorbium, Ferrum, Iodine, Kali, Lauro-Cerasus, Mezereon, Acid. Muriatic., Natrum, Nux Moschata, Plumbum, Psoricum, Ranunculus, Sarsaparilla, Senega, Senna, Silicea, Spigelia, Valerian.

A practitioner, however, must necessarily be consulted where colic is purely symptomatic, in which case it would be highly imprudent for the patient to rely upon his own resources.

Congestion of Blood in the Lower Part of the Abdomen.

This disease is distinguished by a painful sensation of weight, hardness, tension, and heat, in the lower region of the abdomen. When it is of

long standing, it requires the assistance of the physician, and the antipsoric treatment is often necessary to its removal.

When of recent occurrence, and not originating in chronic miasma, but arising merely from errors of diet, it may be readily combated by regimen, by daily and moderate exercise in the open air, and by abstinence from spirits. The patient may, at the same time, have recourse to *Nux Vomica*, of the 24th or 30th attenuation, two globules of which dissolved in six spoonfuls of water, and a spoonful taken every night on going to bed; such dose being repeated successively, or after some days' intermission, will speedily check the disease. Should the remedy fail, the advice of the physician will be indispensable.

Mercury is a valuable medicine in venous congestion of the stomach and the lower part of the abdomen.

Congestion of Blood in the Chest.

This disease attacks young persons of plethoric constitution, and more especially those who are liable to chest affections. Weight and pressure at the chest, palpitations of the heart, asthma, sighings, anxiety, &c. are the most usual symptoms.

Such are frequently the avant-couriers of hæmoptysis, or of some other disease of the lungs, or of the heart. *Nux Vomica* will frequently triumph over these symptoms.

Belladonna is suitable where the asthma is severe, and is accompanied by a constant heaving cough, depriving the patient of rest.

Aurum is a very powerful remedy in cases where the palpitations of the heart are extremely violent.

China is successful in similar cases.

Bryonia is effective against violent and periodical congestions of the chest, accompanied by determination of blood to the head.

Pulsatilla is very powerful against congestion in the head and chest, in connexion with the following additional symptoms:

Inclination to shed tears; irritability; disturbed repose; throbbing and beating, especially in the left side of the head; black spots before the left eye; mucous taste in the mouth; tongue white and moist; great dryness, without thirst; difficulty of deglutition; giddiness, especially in stooping; expectoration of mucus in the evening; palpitations of the heart; tearing pains in the sacrum, on moving; pricking sensation in the arms; empty risings; bad appetite; transitory heats; great debility.

Congestion of Blood in the Head.

This disease more particularly affects persons leading too sedentary a life. It frequently results from natural predisposition, from mode of living, the use of spirituous liquors, the habit of wearing too warm a head-dress. Infancy and old age are the periods of life chiefly predisposed to this affection. The symptoms are generally the following:

Redness and turgid appearance of the face; swelling of the veins of the head; strong arterial pulsations; giddiness; pains in the head, increased by stooping; sleep, broken and disturbed by frightful dreams.

Aconite is, in the first instance, the most appropriate remedy.

Nux Vomica is indicated after the exhibition of aconite, and principally where the congestion arises from a sedentary life, intense mental application, the use of coffee, and of spirituous liquors.

Belladonna is more particularly suitable where we find semi-lateral cephalalgia, violent lancinating and burning, attended with intolerance of light. The administration of this remedy should be preceded by one or more doses of aconite.

It is suitable, again, where, during the cerebral congestion, sparks appear to dart before the eyes, where there is dimness of vision, double sight, hissing in the ears, fainting and drowsiness.

These symptoms exist at the period of dentition, so frequently accompanied by convulsive fits. At the age of puberty, when menstruation is not yet regularly established, this state of things is frequently observable; and indeed, in women generally, where the menstrual derangements have been caused by coldness of the feet, or by reason of suppression from any other circumstance. In such cases we must give aconite, and after one or two doses of this remedy, we may have recourse to belladonna.

In cases where fright and anger have been the exciting cause of cerebral congestion, aconite is the specific *par excellence*.

Where cerebral congestion has been the result of a heavy fall, or severe contusion, we may relieve it by the exhibition of *Arnica*, taken internally, and applied externally as a lotion to the head.

Concussion of the brain is recognizable by the following symptoms:

Giddiness; vertigo; sense of cold in a very circumscribed portion of the head; pressure

over the eyes, which have a tendency to close; disposition to be frightened; vomitings.

In cases where the administration of *arnica* produces bleeding from the nose, we should not interfere with it: the hemorrhage will cure itself.

Dulcamara may be given when there is continual buzzing in the ears, with hardness of hearing.

Coffea is very valuable in cases of congestion occasioned by joy of too lively a character.

Chamomilla in cases of congestion induced by vexation.

Ignatia in cases resulting from concentrated grief.

Nux Vomica in those presenting themselves after a fit of anger.

Opium against those originating in fright.

Crocus in cerebral congestion either previously or subsequently to the menstrual period.

Constipation.

This disease is one of very frequent occurrence in England, and is oftentimes symptomatic of a complicated affection of the digestive organs. In these cases, therefore, the assistance of the medical man is indispensable; and it would be useless to apply one's self to the constipation merely, since the treatment should embrace the malady

in all its bearings. The homœopathic practitioner, who would direct his attention to this symptom merely, would be equally culpable with the allopathic physician who prescribes purgatives for every variety of constipation with which he may have to deal. When the constipation continues during several days, without being connected with any other circumstance, it is frequently the result of too scanty a secretion of the juices essential to digestion, or from the weakness of the intestinal canal. It affects, for the most part, persons leading too sedentary a life, women, and aged men. Diet of too dry a character, a state of pregnancy, the abuse of enemata and of purgatives, are the chief predisposing causes.

The treatment must have reference to the *cause* of the ailment. *Opium* is very salutary in cases of simple constipation, not following in the wake of a long-continued illness, nor resulting from chronic miasma, but which originates in debility of the nervous system, and presents the following symptoms :

Desire to go to the closet, with a feeling as though the passage leading to the rectum were closed ; an impossibility of voiding the fæces ; throbbings in the lower part of the abdomen ; pressure at the stomach ; loss of appetite ; thirst ; dryness in the mouth.

Nux Vomica proves a valuable remedy where constipation is the result of indigestible food, or food taken in too large a quantity; where it occurs as a sequel to diarrhœa, checked by remedies ill chosen, or ceasing spontaneously. Or, again, where the following symptoms are observable:

Loss of appetite; disagreeable taste in the mouth; tongue loaded with mucus; distaste; nausea; lancinating pains in different parts of the lower abdominal region, with tension and pressure; cutting pains in the abdomen; heat; fulness of blood in the face; headach in walking about, in lowering the head; pressure in the temples, relieved neither by sitting up nor lying down; pain in the forehead; in the nape of the neck; disturbed repose; oppression; peculiar heat in the lower part of the abdomen.

The constipation, at times very painful, to which pregnant women are liable, will frequently yield to the exhibition of *Nux Vomica*, alternated with *ignatia*, *bryonia*, and *opium*. We must at the same time enjoin walking exercise every day in the open air.

In cases refusing to yield to these various means, we must consult a practitioner, who will find *Sulphur*, *Sepia*, and *Lycopodium* very frequently indicated.

After an accouchement, where the bowels do

not readily perform by the unaided powers of nature, we must employ *Nux Vomica*, *Ignatia*, *Bryonia*, and *Opium*.

Our choice among these different remedies must be regulated by accessory circumstances, and by the moral tendencies of the patient. *Nux Vomica* responds more especially to temperaments which are ardent, hasty, and inclined to anger; *Ignatia* applies to persons who are gentle, amiable, and vivacious, variable in disposition, sometimes lively, sometimes depressed; *Bryonia* is suitable where the disposition receives a powerful impress from the disease; *Opium* is adapted to persons who appear devoid of energy, and whose bodies are in a state of torpor.

The constipation of newly-born infants most frequently originates in their nourishment being ill suited to their age. This more particularly occurs when they are brought up by hand. In cases where the milk of the mother is insufficient, we must apply such additions only as are very light: for instance, the milk of the cow diluted with water, pigeon or chicken broth, water prepared from grated bread, gruel lightly boiled, &c. When the constipation exists despite of such measures, we must employ *lavements* of milk and water.

Nux Vomica may be administered, either in the

first instance, or upon the failure of the enemata.

Pulsatilla is very valuable in infantile constipation, where the following symptoms are evidenced: temperament quiet and placid; taste acrid and bilious; acid risings in the mouth; retchings; mucus in the mouth; inclination to be sick; burning heat in the pharynx; oppressive tension in the lower part of the abdomen; gripings; livid complexion; debility; shiverings.

This medicine is equally beneficial to women in whom the constipation produces pains on one side of the head, when associated with chilliness, an absence of thirst, melancholy, inclination to shed tears, anxiety, and where the moral tendencies of the patient are mild and placid.

Opium is suitable when the abdomen becomes gradually swollen, and where the disease is unattended with inclination to go to the closet.

Bryonia may be administered where the constipation is accompanied with determination of blood to the head, vertigo, pressure of the temples from without inwards; when, upon stooping, the brain feels as though it would force itself through the forehead; and where there is a tendency to epistaxis.

Sulphur, Veratrum, Plumbum, Conium, Cannabis, Sarsaparilla, Psoricum, Graphites, and

Sepia, are very important remedies in constipation. To render their application effective, however, we must carefully examine all the other symptoms attendant upon constipation; and the study of the *Materia Medica* can alone point out that which will be most eligible.

Contusions.

Contusions very quickly occasion, as all persons are aware, a swelling, more or less extensive, of the injured parts. The tumour is painful, not limited in extent; is more or less hard, and accompanied by extravasation of blood, communicating to the part a peculiar hue, red, deep, mottled, blueish, purple, afterwards yellow, and which is more or less obstinate according to the case and the disposition of the patient.

Arnica is a certain specific against this accident. It is exhibited internally, and had recourse to at the same time as an outward application. In the latter, fine linen rags must be dipped into a solution of the tincture of arnica and water, in the proportion of from ten to twenty drops to the ounce. The injured parts must be moistened with the lotion several times during the day.

Generally, one or two doses internally will suffice, provided the malady be treated at the

onset; and it is seldom required to apply it externally more than one or two days. In serious cases, however, a continued application may be necessary; we must then be guided in the repetition by general rules laid down in the chapter devoted to the subject.

Cramps—Convulsions.

In most instances, cramps are symptomatic of some serious malady, and in such cases it is evident that the medical man alone can direct the treatment. In this place we shall merely consider those spasmodic affections which may be transitory.

Young children are frequently attacked with cramp, from trivial causes, and without any other appreciable derangement of health. In such cases cramp usually supervenes during sleep, appears only in the fingers, or displays itself by a slight twitching of the whole body. Here it has its origin in a cold, in worms, or in colic.

Chamomilla will generally succeed in restoring the equilibrium, taking care to administer it in water, and repeating the dose on every repeated attack. *Chamomilla* is efficient where the child exhibits uneasiness, when one cheek is flushed and the other pallid, when the eyes are half

closed, when the patient appears without intelligence, is convulsively affected in the arms and legs, rolls his head uneasily on the pillow, moans, and suffers from thirst.

Ignatia may be administered under the following circumstances: Frequent accession of heats over the body; waking by starts from light slumber, and composing himself again with difficulty; uttering loud cries, and feeling a general tremor; convulsive movements of the muscles, or of the limbs.

Mercury is an appropriate remedy, when, in addition to the convulsive motion of the limbs, we may observe swelling and hardness of the lower belly; ejection of a watery fluid; change in the intellectual faculties; general debility; and where all these symptoms are evidently referrible to the presence of worms.

Women of delicate and irritable temperament are attacked previously to their monthly periods, or after a cold, or from some moral disturbance, with spasmodic and constrictive pains in the lower part of the abdomen; and even with convulsive movements, more or less violent in the limbs, the pains contracting and pressive, and extending from the loins to the pelvis and the thighs.

Coffea is an excellent remedy in such cases, particularly where its use has not been habitual.

If the patient have been accustomed to this beverage, and it fail in its application, we must have recourse to *Pulsatilla*; especially if there be shivering, inclination to weep, anxiety, and when the symptoms become exacerbated during the night. *Chamomilla* is better adapted to those persons whose vivacity, quickness, and extreme susceptibility of suffering, constitute their moral characteristics.

Ignatia is efficient in hysterical affections, and occasions the disappearance of cramp when in connexion with such symptoms as the following:

Nausea; lipothymy; loss of sense; general chilliness; pallor; increase of sensibility to light or noise; painful tumefaction of the lower abdominal region; twisting; digging pain in the lower part of the abdomen; increase of heat in the body; pulse sometimes rapid and full, at others weak and small.

Opium will relieve spasmodic affections induced by fright, and which present, as accessories, loss of intelligence, a gloomy expression of countenance, general tremor, diarrhœa, difficulty of respiration, weakness in all the limbs.

Men of delicate fibre occasionally experience cramp-like sensations, such as impeded respiration, yawnings, constriction, oppression about the chest, general laxity, and shiverings.

Camphor, applied from time to time to the nose, will cause these symptoms to retire.

If these symptoms are evidenced by persons of violent and hasty temperament, and who give way to tears, complaints, and abuse, *Nux Vomica* will be the suitable remedy.

Ipecacuanha is serviceable in convulsions, with great oppression at the chest, with distaste for everything but water, with nausea, vomitings, and diarrhœa.

Lemon-juice, *Cicuta*, *Causticum*, *Cannabis*, *Aurum*, *Arnica*, *Veratrum*, *Stramonium*, *Spigelia*, *Moschus*, *Menyanthes*, *Aconitum*, *Arsenic*, *Bel-ladonna*, *Bryonia*, *Cocculus*, *Cuprum*, *Hyoscy-amus*, *Angustura*, *Lachesis*, may be employed in various forms of convulsion.

Cramps of the Stomach.

This malady is one of frequent occurrence; appears to be sometimes hereditary, to come on in paroxysms, and may continue for years.

At first, there is merely pressure, with tightness and pinching more or less severe, affecting the pit of the stomach, and the left side of that viscus. The pain frequently extends to the back, where strong pressure is felt as from a hard

substance. The pains afterwards become very perceptible, being hot, cramp-like, and insufferable, extending themselves along the œsophagus, with a sense of strangulation. When at their height, they are very severe, accompanied by constriction, twistings, anxiety, pressure, shootings in the breast and in the back, colic, increase of saliva, sobs, and lipothymy. The fit continues at first for some minutes, afterwards for a quarter of an hour, an hour, and even half the day, and terminates in vomiting. The pain usually disappears when the stomach is empty, or contains merely light and soft food, and again supervenes after a repast. It is increased when standing upright; whilst a forward or bent position, and frequently even pressure, will procure relief. Sometimes the region of the stomach is a little hard and inflated, more usually contracted, and very rarely sensitive or tense. We frequently remark an inclination to be sick, but the inclination is not always responded to. When the latter symptom presents itself, a brief melioration is the result.

Cramps of the stomach are more or less obstinate according to the circumstances to which they are referrible, and frequently resist for a long time the most judiciously directed

treatment. At times they disappear for several weeks, and return at fixed or irregular periods.

The most usual causes of this malady, are errors of diet, frequently and for a long time repeated; irregularity in eating and drinking; the inordinate use of spirituous drinks and coffee; grief, vexation, frequent colds, particularly in the feet, &c. All these causes, however, are often merely occasional and exciting; and cramps in the stomach originate in some chronic miasma, which develops itself under the influence of the causes alluded to above. In such cases we must institute an antispasmodic treatment, methodical and prolonged, in order to triumph over the disease.

When the affection is attributable to the causes above mentioned, it is generally to be cured by the following remedies: *Nux*, *ignatia*, *pulsatilla*, *coffea cruda*, *chamomilla*, and *cocculus*.

Nux Vomica is especially applicable to persons who are addicted to coffee and spirituous liquors, more particularly when we detect the following symptoms:

Constriction, pressure, pinching, and tearing; cramps in the stomach, with a feeling as though the clothes compressed the body in the epigastric region; a sensation which may be compared to an accumulation of wind in the left side, be-

neath the ribs; constriction of the chest, which in many instances extends itself as far as the shoulders and the loins; nausea, particularly during the fit; accumulation of water in the mouth; rising in the mouth of sour and bitter fluid, with or without a sense of burning in the pharynx; palpitation of the heart, with anxiety; sour and putrid taste in the mouth; constipation; tumefaction of the lower part of the abdomen from flatus; at times semi-lateral headach; at other times, pressive pains in the forehead.

When the nervous system is highly excitable, the sensibility increased, the disposition of the patient very irritable, and every circumstance annoys him, we shall derive much advantage from chamomilla; more particularly where there is pressure in the region of the stomach, painful swelling at the pit of the stomach, and on the left side; asthma, anxiety, and a pressive pain at the vertex of the head.

It may be remarked that a little coffee generally assuages this species of cramp, while it serves to aggravate that which yields to the application of *nux vomica*.

When the disease has originated in the abuse of chamomile tea, this remedy will not meet the case: it is then that *ignatia*, *pulsatilla*, and *coffea* will be appropriate. *Coffea* is especially indicated in

cramps which appear insupportable to the patient, by reason of an increase of sensibility. Cramps in the stomach, excited by an abuse of coffee, are remedied by the exhibition of chamomilla.

When the cramps are connected in their paroxysms with nausea and vomiting, with severe darting pain in the pit of the stomach, they will yield to the administration of *Ipecacuanha*.

Cocculus may be administered after the exhibition of nux, should the latter fail in the entire removal of the disease; or, again, when super-added to the symptoms which characterize the employment of nux vomica, we have constipation and pain in the lower part of the abdomen, relieved by the discharge of flatus; when there is an increased flow of saliva from the mouth; and when the disposition of the patient is fretful and sombre, and he seems as though he were wrapped up within himself.

The following medicaments are valuable in certain descriptions of cramps in the stomach :

Belladonna, Stannum, Staphysagria, Bryonia, China, Magnesia, Bismuth, Arsenicum, Plumbum, Argilla, Carbo Vegetabilis, Calcarea Carbonica, Barytes Carbonica, Lycopodium, Sepia, Petroleum, Atriplex Olida, Carbo Animalis, Phosphorus, Spiritus Nitri Dulcis, Sulphur.

Cries of Young Infants.

The violent crying of very young children, continuing frequently during one or many weeks, without apparent cause, and almost without intermission, will frequently yield to the exhibition of an extremely minute dose of belladonna, chamomilla, or coffea. This latter substance is particularly appropriate where the nurse is not habituated to coffee.

Chamomilla is applicable when, in addition to the cries, there is much agitation, and the child is very restless.

Jalap is suitable in cases unattended with diarrhœa.

Senna, in cases of flatulence and disturbance of the circulation.

Rheum, where there is a disposition to relieve the bowels in the form of colic.

The best mode of exhibiting medicines to very young children is to dissolve a globule in 20 teaspoonfuls of pure water, and to give a spoonful to the little patient every quarter of an hour, every half hour, or every hour, according to the effect derived from the medicine.

Milk-crusts.

This disease principally affects infants at the breast. It consists of vesicles filled with clear and transparent lymph, forming clusters, and covering a surface sometimes considerable in extent. The pimples usually make their first appearance on the face, and will sometimes extend themselves over the entire surface of the body. They soon become yellow, break, and subsequently form very hard crusts. Should the part where the milk-crusts have their seat be swollen, if the skin be very red and tender, and if the children be extremely restless, and endeavour to scratch themselves, we must administer one or more doses of *Aconite*, succeeded, after an interval of thirty-six or forty-eight hours, by the *Viola Tricolor*. Two or three globules of this medicine are dissolved in water, and administered during several days, according to circumstances; and the same medicine may be repeated once or many times.

This treatment readily meliorates the condition of the child; sometimes, however, it requires weeks to surmount the evil.

Should this malady originate in psora, transmitted by the parents, we shall not at all times succeed in subduing it by the two medicines

alluded to. In such cases we must have recourse to antipsoric remedies.

Sulphur takes precedence after the exhibition of the *viola tricolor*, one or two doses being sufficient to effect a perfect cure.

Arsenic, Barytes, *Lycopodium*, *Carbo Vegetabilis*, *Causticum*, *Bryonia*, *Dulcamara*, *Graphites*, *Hepar Sulphuris*, *Psoricum*, *Rhus*, *Sarsaparilla*, and *Sepia*, are medicines frequently of great service in the more obstinate cases. It is for the physician to determine which may be preferable, and he will arrive at this choice by a perfect knowledge of their respective properties.

Itchings.

This disease is frequently obstinate and very distressing, and chiefly attacks persons who are declining in years. It has its immediate origin in slight nodosities, colourless, scarcely perceptible, soft, smooth, isolated, without vesicles, desquamation, or eruption, properly so called.

The scratching, however, produces a slight redness, with discharge of lymph, clear or sanious, visible at the top of the nodosities. This assumes the appearance of a number of small humid points, which form, in the course of twelve or twenty

hours, as many minute crusts or scales, about the size of a pin's head, white, yellowish, or dusky, according to the character of the fluid, and frequently rendering the skin rough to the touch. The scratching at times, too, produces vesicles. When this affection is of long standing, it frequently degenerates into a well-defined eruption, and does not yield to time. The scratching, so far from relieving the pain, frequently adds to its violence, and causes much inflammation and heat. The annoyance increases towards evening, on exposure to warmth, after having recourse to stimulating food and beverage, by motion, and during the monthly periods. Occasionally it presents a slight intermission, chiefly during repose or occupation; it even disappears altogether, to return at fixed or variable periods, especially at changes of temperature, spring, autumn, &c. These itchings are more usually experienced behind the shoulders, at the nape of the neck, in the bosom, at the inner part of the thighs, and but seldom between the fingers. Sometimes the whole body participates in the inconvenience.

They originate, in certain cases, in absence of cleanliness, particularly among the lower classes; in checked perspiration; in plethora; in the use, either as food or as beverage, of substances which

are sweet, poisonous, aromatic, or spirituous; affections of the liver, gout, scurvy, and above all in psora, the last of which may even be considered as the originating cause in a majority of cases, the others being regarded in the light of excitants.

The remedy most certain and most prompt is *Sulphur*, which may be administered in solution with water, in the proportion of from one to three globules every week, and repeated according to circumstances. We must be careful not to exhibit a second dose whilst the action of the first is in progress.

If the itching be diffused, pricking, and burning, and assume the appearance of flea-bites, is unaccompanied by any visible mark, is felt chiefly in the evening on going to bed, if it give way upon the slightest scratching to reappear in another part, we must order the *Ignatia amara*, in the same doses as the sulphur.

When this treatment, united with habits of cleanliness, may prove insufficient to check the disease, and particularly when connected with a complication of symptoms, we must profit by the advice of a medical practitioner.

Dentition.

Dentition is rarely observable in young children without bringing in its train symptoms more or less serious, and which the physician is called upon to remedy.

The soft and gelatinous substance of which, at a later period, the teeth are formed, is enveloped in a membranous sac, and secreted within the bone of the jaw. This substance gradually increases, and, from its consistency, opens for itself a passage by causing a dissolvent irritation in the bone. The dissolved portion is removed by the absorbents, and it only remains to pierce the gum, which by degrees becomes inflamed and swollen. The gum being but slightly sensitive, the pain is generally trifling at the point of perforation.

When dentition advances without unfavorable symptoms, it requires no assistance from art. If the swelling of the gums, however, be considerable—if they are red, hot, and painful—if the child be incessantly drivelling—if he frequently introduce his fingers into his mouth—if he forcibly bite the nipple when at the breast, or suddenly let go his hold—and if he be restless, we must oppose this

irritation by aconite; one globule dissolved in four spoonfuls of water, and a spoonful given every two hours.

When the child is as it were delirious, is unwilling to be pacified, and unable to sleep, we must have recourse to *Coffea*, exhibiting one or two globules in the four and twenty hours.

If the mother make habitual use of coffee, we must then apply to *Chamomilla*, administered in the same manner.

When dentition is accompanied by a dry cough, with panting, or hurried respiration, chamomilla is very serviceable; should it fail, however, *Belladonna* will succeed in restoring the health of the patient.

These medicines may be employed under other circumstances similar in character to those mentioned above: they are especially efficacious where the child has for several nights experienced restlessness, burning heat, constant agitation, ardent thirst, redness of the skin, tremor of the limbs, anxiety, sobbings, sighs; respiration short, quick, and loud; evident oppression at the chest, redness of the eyes; startings, isolated, and recurring at intervals, and even convulsion of the limbs. What should influence our choice in favour of belladonna, would be evident and well-defined symptoms of cerebral congestion.

Chamomilla is particularly suitable where the general symptoms are the same; and where the congestion appears to influence the abdomen, accompanied by gripings, diarrhœa, and green evacuations.

In cases where the diarrhœa is watery, and very profuse, we may oppose it with *Mercury*, one globule of the twelfth attenuation being given in the course of twenty-four hours. The diarrhœa, however, being favorable to dentition, we must interfere with those cases only in which the symptoms are violent. Constipation, on the other hand, should be carefully obviated with *Nux Vomica*, which will, in most instances, readily meet the case.

Ignatia is appropriate in cases of convulsions which have withstood the action of *Belladonna* and *Chamomilla*. We may repeat the *ignatia* according to the severity of the attack; one dose every quarter or every half hour.

Cina, *Rhus*, *Arsenic*, *Arnica*, *Hyoscyamus*, *Ammonium*, *Cicuta*, *Stannum*, *Cuprum*, and *Pulsatilla*, are likewise valuable in derangements resulting from dentition.

Passion.

We frequently remark that a violent fit of passion will exercise a prejudicial influence upon the health, and give rise to diseases more or less important. After the first disturbance induced by anger, we may administer a dose of *Chamomilla*, should the following phenomena present themselves:

General excitement, agitation, tremor, heat, and redness of the face and eyes; compressive pain in the head, palpitation of the heart, bitter taste in the mouth, excess of saliva, loss of appetite, constriction from tightness of the stomach, nausea, vomitings of green and bitter substance, gripings, diarrhœa, and thirst. In such cases *Chamomilla* is very appropriate.

Ignatia is indicated for those persons whose passions are concentrated within themselves.

Nux Vomica is appropriate in very nearly the same cases as *Chamomilla*, more especially when the patient indulges in complaint, murmuring, and tears.

Pulsatilla is applicable to persons of acute sensibility, of irritable temperament, delicacy of fibre, meekness of disposition, inclination to shed tears, paleness of countenance, shivering, headach,

absence of hunger and thirst, bitterness in the mouth; diarrhœa, glairy or green, occurring chiefly in the night; continual rumblings in the abdomen, alternations of heat and cold, and vigilance occasioned by an internal anxious heat.

When a violent paroxysm of anger has thrown the patient into a state of extreme distress, so that he scarcely knows which way to turn himself, *Arsenic* will be the most appropriate remedy.

When dread is superadded to passion, *Aconite* becomes a specific.

Should this remedy, however, fail in the removal of all those symptoms, we shall be able to select one, from those enumerated above, which will perfect the cure.

Derangement of the Stomach.

This indisposition originates in excess of nutriment, or from the food and drinks being heavy, indigestible, gross, acid, flatulent, tainted, &c. The symptoms of visceral derangement generally declare themselves very quickly after a repast, and are as follows :

Risings in the mouth, of an acid character, or tasting of the food which has been eaten ; disrelish,

weight, pressure and pain in the stomach; swelling and fulness of this viscus; quivering of the lower lip; vertigo; heaviness in the head; tickling or scraping in the throat; expectoration; hemoptysis; unpleasant taste in the mouth; loaded tongue; inclination to vomit; heat in the throat; sensation of weakness; anxiety; face puffed up, pale, covered at times with various coloured spots; discharge of fetid wind; the urine sometimes admixed with mucus, &c.

These symptoms, which usually supervene some hours after the repast, soon terminate in acid vomitings participating in the nature of the food, and sometimes presenting a greenish appearance. There remains, however, for some days, loss of appetite, pains in the abdomen, or diarrhœa.

Persons of delicate constitution, and subject to cramps in the stomach, suffer much from indigestion and its concomitant ills; among which may be mentioned paroxysms of fever. Occasionally the food undigested, or taken in too large a quantity, causes violent symptoms, such as might result from poisoning, &c.; but they are generally brief in duration.

Arnica may be administered where the derangement of the stomach is referrible to a diffused and continued excitement of the nervous system occasioned by intensity of thought, late hours, excess

of venery, bodily fatigue, and when accompanied by the following symptoms :

Vertigo ; headach, especially above the orbits ; giddiness ; heat of the head ; dryness of the tongue ; putrescent, sour, or bitter taste in the mouth ; unpleasant odour in the mouth ; distaste for smoking of tobacco ; want of appetite ; desire for acid food ; tongue loaded with a yellowish mucus ; risings in the mouth, and tasting of the food ; fulness in the pit of the stomach ; nausea ; inclination to vomit ; heaving of the stomach ; flatulence, colic, swelling of the abdomen ; uneasiness and debility ; heaviness in the limbs ; disagreeable heat ; cracking of the knees ; disturbed slumber ; frequent wakings ; waking suddenly ; anxious and distressing dreams.

Chamomilla and *Nux Vomica* will succeed in such cases, provided there be no circumstance to contra-indicate their use.

Nux Vomica will check derangements of the stomach produced by nocturnal debauch, abuse of wine or coffee, or by a cold, and when the patient displays the following symptoms :

Sensation of giddiness ; obnubilation of the head ; sensible weight in the occiput ; drawing pains in the teeth, sometimes in one of the superior, sometimes in one of the inferior molars ; tingling in the ears ; heat in the face ; efflorescence in the forehead, red and isolated ; the same

in the nose and at the angles of the mouth; tongue white and loaded; dryness of the mouth, without thirst; collection of sordes in the mouth; heat in the throat; failure of appetite; loss of taste; nausea; accumulation of water in the mouth; vomitings; colic; pressure at the stomach; tension in the lower region of the abdomen; evacuations scanty; constipation; disturbed sleep; disinclination for intellectual pursuits; great relaxation; twitchings in the limbs; discontent, querulousness, and inward agitation.

Derangement of the stomach, occasioned by vexation preceding a repast, presents the following symptoms:

Heat and redness of the face; painful sensation in the head; redness of the eyes, with burning heat; excess of excitement pervading the nervous system; great sensibility; failure of appetite; continuous bitter taste in the mouth; bilious eructations; vomitings of green and bilious substances; colics; prostration; disturbed slumber; perpetual agitation; frequent wakings.

Chamomilla will generally succeed in dispersing all these symptoms. Should uneasiness, however, in the epigastrium continue, the appetite still fail, the patient feel great lassitude, with absence of thirst and loaded tongue, *Pulsatilla* will be the

remedy to confirm the equilibrium of the functions.

Nux Vomica, on the other hand, may be prescribed with success, where we find constipation, and the temperament of the patient violent, with an inclination to quarrel.

Bryonia, again, may be employed under similar circumstances, especially if the patient be costive, feel shiverings, and it is perceived that his moral economy has undergone a signal change under the influence of anger.

Pulsatilla is more especially applicable where the derangement has resulted from the food being gross and flatulent, and when the under-written symptoms exhibit themselves :

Bitter and salacious taste in the mouth ; taste resembling that of putrescent fish or tallow ; mucus in the mouth ; scraping in the throat ; bilious eructations ; loss of appetite ; aversion to warm food ; fulness of the stomach ; swelling of the lower region of the abdomen ; tension beneath the false ribs ; borborygmus ; costiveness, or heavy, difficult, and scanty evacuation in the morning ; chilliness ; prostration ; twitchings in the limbs ; ill humour, vexation.

In the vernal and autumnal seasons, among persons who reside in humid situations, in the

vicinity of stagnant water, in marshy districts, and in closely-confined or densely-populated apartments, we observe a peculiar gastric derangement, presenting the following characteristics :

Loss of appetite, without unpleasant taste ; constant satiety ; heaviness and fulness of the abdomen ; constriction, prostration, and debility ; vertigo ; deafening headach ; flushes, occasionally alternated with chills ; urine thick and clouded ; sensibility to external impressions ; disturbed repose ; ill-humour.

China, in one or more doses, speedily arrests this train of symptoms, the not unusual precursors of Intermittent Fever.

Diarrhœa.

Diarrhœa is an evacuation, more or less frequent, resulting from disease of the intestinal canal. The voided substance is at times copious, diversified in character, and with or without admixture of fecal matter.

Occasionally this affection is very trivial, and, in consequence, is attended with little danger ; at other times it assumes a graver aspect, more especially when injudiciously treated.

The most frequent causes of this malady are

colds, errors of regimen, &c. The concomitant symptoms are loss of appetite, more or less defined; pain in the stomach; swelling and tension of the lower part of the abdomen; borborygmus; dryness and coldness of the skin; thirst, diminution of urine; prostration; intermittence and irregularity of the arterial pulsations; occasional pains in the abdomen. Then, again, we have cutting and tearing pains, sometimes in one part, sometimes in another, with inclinations to vomit, subsiding for the moment to return after a brief interval. Severe and long-continued diarrhœa occasions burning pain in the rectum, tenesmus, debility, alteration of feature, and even lipothymy.

Warmth, in conjunction with abstinence and a little coffee, is sufficient to remedy slight attacks. Those cases, however, which have been produced by colds, must be treated with china, provided the motions are frequent, liquid, sour, clear, and without admixture of fecal matter; are preceded by violent colic, spasmodic and pressive pain in the bowels, and accompanied by pains in the abdomen; eructation; borborygmus in the abdomen, with sense of debility in the same organ.

Chamomilla is suitable when the pains in the abdomen are violent and tearing, afford no relief to the patient, and oblige him to writhe and run about here and there; when there is a feeling as

though the abdomen were entirely hollow, with constant movement in the bowels, or as though a ball were rolling about in the side; when the eyes are encircled by a blueish ring; nausea; vomiting; and the motions mucous and watery, having the odour of rotten eggs.

Chamomilla, again, is successfully employed against diarrhœa originating in anger, and where there exists a bitter taste in the mouth; risings; a sense of emptiness or repletion in the stomach; pressive pain in the head; general debility; repeated evacuation of matters, which are green, aqueous, hot, and very fetid.

Mercury is suitable where the cold has been produced by exposure to night air, and there exists a pinching pain in the hollow of the stomach; borborygmus; a feeling of relaxation as though an evacuation would take place; cutting pain in the lower bowels, accompanied by pressure; tearing pain in the lower part of the abdomen, which feels very cold; nausea; aversion; titubations; shiverings; sudden and frequent desire to relieve the bowels, in the first instance without success, but succeeded by dejections, frequent, profuse, watery, and green, with disposition to faintings.

Pulsatilla is of great service in cases of diarrhœa proceeding from errors of diet, and presenting

such indications as loss of appetite, with sensation of dryness and slight burning of the tongue; absence of thirst; a salt and bitter taste in the mouth, without the food being unpleasant; risings in the mouth, as of rotten eggs; hiccough; nausea; disposition to vomit; vomitings; fulness of the lower region of the abdomen; colics; much flatulence; rumbling in the intestines; pressing desire to go to stool; restlessness; frequent evacuations.

Opium will quickly remedy diarrhœa, occasioned by fright, anxiety, powerful emotion, &c.

Dulcamara is particularly efficient in aqueous diarrhœa, induced by cold, and accompanied by neither pain in the abdomen nor by any other accessory indisposition.

Dulcamara is valuable again in cases of infantile diarrhœa, in teething, and which is usually attended with fever and loss of appetite.

Sulphur is an important remedy in diarrhœa among children of psoric habit, and where the following symptoms are evidenced:

Acrid diarrhœa, excoriating the anus and surrounding parts, and even the thighs, and exciting in such parts a miliary eruption, occupying a greater or less extent of surface; gradual emaciation, and consumption arriving at its last stage.

Pains following Childbirth.

Notwithstanding these pains rarely constitute disease, properly so called, it may be of service to point out the means of relief or dispersion. They may now and then prove sufficiently intense to induce consequences more or less serious.

These pains are analogous to those preceding the labour; they extend from the lumbar region to the pelvis or the sacrum, are accompanied by irritation in the rectum and the bladder, and affect the pubic region and the posterior portion of the lower part of the abdomen.

They succeed delivery even under the most favorable auspices; come on in paroxysms day after day; are unconnected with fever or abnormal discharge; and during the intervals, howsoever brief, the patient is in perfect health, and her appetite and sleep return. These pains rarely occasion any change in the lochia, the secretion of milk, or the alvine evacuations, and they are terminated occasionally by a discharge of blood. Pressure does not increase them, neither is the abdomen swollen nor tender. These after-pains are at times more distressing than those of labour; and it has been remarked that they are generally

more severe in women who have borne several children. If trivial in character, they require but little interference, nature herself restoring the equilibrium. Should they become serious, we may oppose them with *Chamomilla*, which, if it fail in their entire removal, will very greatly relieve. Should the pain prove very severe, and cause the patient to feel as though the intestines were cut into pieces, we may derive beneficial results from *Coffea*.

Arnica has been employed most successfully, in the form of lotion to the organs of generation; and the exhibition of a dose or two, internally, will contribute in a singular degree to a favorable termination.

In all cases of contusion, the *Arnica* proves invaluable; it must, therefore, of necessity, do good when employed subsequently to an accouchement. If this remedy do not altogether prevent the accession of after-pains, it will, in most instances, produce a modification; while *Chamomilla* will accomplish the remainder.

Poisoning.

Accidents of this nature claim the unremitting energies of the most skilful practitioners. It is

nevertheless important, that the public should be put in possession of the means by which to recognize such cases, and afford the earlier means of relief, while awaiting the arrival of competent assistance.

Where violent symptoms are suddenly evidenced, and without appreciable cause, in persons of sound health or but slightly indisposed, we may infer that such phenomena result from the exhibition of poison. Poisons are either narcotic or corrosive. The first division, belonging chiefly to the vegetable kingdom, comprises *Opium*, *Belladonna*, *Datura Stramonium*, *Hyoscyamus*, *Cicuta*, *Secale Cornutum*, certain species of mushroom, &c. The bitter almond and the laurel leaf are poisonous; and being introduced into the stomach in too large a quantity, may become prejudicial to health; while the prussic acid, extracted from the leaves, is still more to be dreaded. Under the same head, we may class such atmospheres and gases as are obnoxious to respiration; suffocating vapours, such as carbonic acid gas, the emanations of sulphur and charcoal, air pent up and vitiated by the breath of too large a number of persons, the odour arising from an assemblage of flowers in a room, &c.

These poisons immediately give rise to the following phenomena:

Painful weight and pressure in the stomach; anxiety; aversion; vomiting; somnolency; confusion in the head; disturbance of the intellectual faculties; vertigo; dulness; coma, &c. The patient feels weak, mutters incoherently, totters, his pulse is variable, generally slow and soft, sometimes strong and rapid. In addition to these symptoms, we find giddiness and stupor, breathing laboured and stertorous, buffoonery, delirium, convulsions, paralysis, and lethargy.

The corrosive poisons are chiefly derived from the mineral kingdom: these are arsenic; verdigris, or the green oxyde of copper; mercury, with its different preparations; very strong mineral acids, such as sulphuric, nitric, and muriatic acids in their concentrated form; certain plants whose juice is acrid and corrosive; among others, for instance, the *Daphne Mezereon*; vegetables uniting a corrosive quality with a narcotic principle, such as *Digitalis* or *Aconite*. *Cantharides* also, on being introduced into the stomach, corrode with singular virulence.

The first results from corrosive poisons are as follows:

Acrid, burning, and very painful sensation in the throat, pharynx, and stomach; pinching pain at the pit of the stomach; inexpressible anxiety; nausea and vomitings; vomiting of matters tinged

with blood; sometimes violent diarrhœa, at others, on the contrary, costiveness, with swelling of the abdomen; abdominal pains, cutting and extremely acute.

In the first instance, while awaiting the arrival of the medical man, vomiting should be encouraged by drinking abundantly of warm water, or, what is still better, water in which a quantity of butter has been dissolved, greasy milk, oil, &c. Patients should be encouraged to drink as much as possible, since the removal of the poisonous principle is apportioned to the amount of vomiting. Persons who vomit with difficulty, should provoke it by tickling the pharynx with a feather or the finger, or by drinking abundantly of strong coffee.

If the poison be vegetable, after having evacuated the stomach by vomiting, the most efficient remedy will be spirits of camphor, of which we may give the patient a drop every three minutes, or less frequently, in proportion as the symptoms subside.

Swelling of the Cheeks.

When this affection is the result of toothach, it usually subsides with the cessation of the exciting cause, and without having recourse to any particular remedy. Should the swelling continue,

however, we must enforce some specific treatment : this will, in most instances, consist of the antidote of the remedy that has relieved the toothach : as for instance, *Pulsatilla*, if the primitive affection have been removed by *Mercury* or *Chamomilla*, and so on of the rest.

If the tumefaction present much obstinacy and hardness, our most effectual remedy will, in most instances, be the 6th attenuation of *Arnica*.

Mercury is particularly efficient where the swelling is accompanied by erysipelatous inflammation, tearing pains, and profuse salivation.

Should the inflammation be violent, and the remedy be adopted too late, the general result is suppuration. There is formed upon the gums, or upon the neighbouring parts within the mouth, a small, hard, and circumscribed tumour, which, as the general swelling is reduced, enlarges, becomes gradually soft, bursts, at the latest, after a few days, and discharges a considerable quantity of matter. The formation of pus may be accelerated by the application of figs boiled in milk. When the suppuration is established, we must leave the cure in the hands of nature. Warm poultices of linseed-meal will relieve the patient, and are in perfect accordance with homœopathic treatment.

*Enlargement of the Glands of the Neck
and of the Ears.*

The glands of the ear, and those beneath the chin, occasionally become swollen, as a sequel to toothach, a cold, &c. A disposition to toothach is very generally succeeded by an enlargement of the glands, and we are frequently obliged to combat this glandular affection, in order to prevent suppuration, induration, &c.

Mercury, in such cases, is a valuable agent, and may be repeated several times, if necessary.

Hepar Sulphuris is of great service when the pains have subsided, or have altered in character, and produce throbbing in the place of lancinations and distention.

If, notwithstanding the exhibition of suitable remedies, the glands attain considerable hardness, *Dulcamara*, taken at intervals, will afford much relief.

Chilblains.

Chilblains, as all persons are aware, are specific inflammations of the skin, resulting from cold. The hands and feet, and especially the great toes, are more frequently affected than any

other part. This inflammatory affection usually displays itself at the commencement of winter, and during variations of the temperature. A single remedy will rarely suffice for its removal, and we are frequently compelled to resort to many.

Chamomilla and *Arnica* are generally entitled to the preference. Considerable inflammation, throbbing pains, tumefaction, and a reddish-blue appearance of the affected part, require the administration of *Pulsatilla* and of *Belladonna*. If the chilblain be of long standing, we may give *Sulphur*, *Nitric Acid*, or *Petroleum*.

Hoarseness.

This affection is distinguished by harshness of tone, indistinct enunciation, crying, cough, &c., and produces, in itself, neither pain, asthma, nor wheezing. It is the result, frequently, of cold, checked perspiration, &c., and is generally accompanied by a dry cough, or by the expectoration of a watery or saltish substance, soon becoming mucous, and more or less profuse and viscid. The hoarseness is scarcely perceptible during inspiration, but is more decidedly

marked in expiration, the act of speaking, coughing, &c.

Chamomilla is applicable where the hoarseness is associated with cold in the head, dryness of the eyelids, collection of mucus in the larynx, sense of burning in the gullet, dryness in the mouth, thirst, dry cough, with prurient irritation in the trachea, febrile excitement in the evening, chilliness, heat, discontent, inward vexation at trifles, gravity, inclination to silence, aversion to lengthened discourse.

Nux Vomica is applicable where there is a smarting in the throat; painful sense of dryness in the larynx, producing cough, accompanied by pain at the pit of the throat; dry, hoarse, and deep cough; cold in the head; painful respiration; vigilance; heat, at night, in the hands and cheeks, with coldness of the feet, and shiverings; accession of chills in the afternoon, succeeded by burning heat and thirst; confusion in the head in the morning; discontent; aversion to occupation; plaintive murmurs; wrangling and disputes upon unimportant matters; obstinate self-will.

Pulsatilla is a valuable remedy in the following train of symptoms:

Hoarseness; hardness, with painful sense of excoriation at the back of the throat; roughness of the palate, with dryness of the throat; pain in

the throat on swallowing; cold in the head, with sanguineous mucosity from the nose; cough, with expectoration and pain in the chest; tickling at the pit of the throat, inducing cough, with evening exacerbation; cold, with twitchings in the limbs; access of fever towards evening; external heat, sense of fatigue, and prostration; internal heat during the night, with skin perfectly dry; disturbed slumber, broken and agitated by dreams; taciturnity, disposition to tears, sensibility, indifference, irresolution, frequent desire for change.

Mercury is often of service after the exhibition of the remedies already pointed out, and will quickly effect a cure.

In this, as well as in all other affections, the repetition of the dose must be regulated by the result.

Belladonna, Bryonia, Calcarea, Carbo Vegetabilis, Causticum, Drosera, Dulcamara, Hepar, Manganum, Petroleum, Phosphorus, Sulphur, Sepia, and Spongia, may be pointed out as serviceable in complicated or chronic cases; and the practitioner will readily select his agent by consulting the *Materia Medica Pura*.

Erysipelas.

Erysipelas is a superficial inflammation of the skin, and is indicated by swelling, redness, and heat of that organ. These symptoms are accompanied by pain, more or less severe, and of a burning character; the finger being pressed upon the painful part leaves a white impress, which soon resumes its red appearance; the hue is of a bright or deep red, sometimes assumes a yellowish cast; the skin glistens, the patient experiences tension and itching, with fever more or less ardent.

Erysipelas chiefly attacks the face and the lower extremities, and especially the joints. Women and elderly persons appear more particularly susceptible.

This affection displays itself chiefly when under the influence of derangement of the digestive functions, powerful emotion, anger, passion, or cold. In other cases, it is evidently the result of some epidemic cause.

Erysipelas will sometimes assume a very serious aspect, and consequently require the most careful attention.

When the case is simply such as we have

described, we are certain to derive very prompt relief from the administration of one or two doses of *Aconite* at the onset, to check the febrile action. This medicine must be followed up by *Belladonna*; or, in the absence of fever, we may begin immediately with the last-mentioned remedy.

Bryonia is of signal benefit where the erysipelas more particularly affects the articulations.

Pulsatilla is applicable where the disease shifts rapidly from one part of the body to another.

In certain cases we perceive vesicles filled with serosity, forming themselves upon the inflamed surface: these are of greater or less dimensions, and resemble small blisters. When these phenomena present themselves, the *Rhus Toxicodendron* should follow the exhibition of *Aconite*.

Under certain forms of erysipelas, and under the guidance of an assemblage of adventitious symptoms, the practitioner will derive much valuable assistance from the following remedies:

Ammonium, Antimonium, Arnica, Arsenicum, Borax, Calcareas, Camphor, Cantharides, Carbo Animalis, Graphites, Hepar, Lycopodium, Mercurius, Nitric Acid, Phosphorus, Ruta, Sulphur.

Quinsy.

Quinsy is an inflammatory affection of the lining membrane of the throat, of the velum palati, and of the tonsils. This affection occasionally assumes a formidable character, and consequently requires very prompt attention. When the disease is imperfectly cured, there results very frequently a chronic enlargement of the tonsils, with elongation of the uvula, which symptoms are to be relieved only by a prolonged and radical treatment. Since in such cases we shall invariably find a psoric tendency, either constitutional or acquired, it is among the antipsoric remedies that the practitioner will find the suitable means of relief.

Chamomilla is an efficient remedy in quinsy, when characterized by dryness of the throat, thirst, difficulty of swallowing, the sensation of a foreign body in the gullet; when there is an inclination, but at the same time an inability, to swallow; where there is throbbing pain, accompanied by fever towards evening, alternations of heat and cold, melancholy temperament, &c.

Ignatia is suitable in quinsy, characterized by the preceding symptoms, and united with the especial sense of feeling, during deglutition, the presence of a ball in the throat, with swelling; and

when, in the act of swallowing, we feel the sensation as of a wound.

Mercury is efficient in this affection of the throat, where the act of deglutition is characterized by a burning sensation in the throat; the throat appears contracted, the pains are lancinating, and extend as far as the ears; the tonsils are swollen and throb during deglutition; the posterior portion of the tongue swollen, as also the gums; unpleasant taste in the mouth; and where, as an indication altogether characteristic, the mouth is filled with saliva, more or less viscid and tenacious; febrile action, more or less apparent towards evening, with alternations of cold and heat.

Two or three doses of mercury are generally required to check this affection.

Pulsatilla is in request where the throat experiences a sense of soreness, and the gullet is dry; when there is cutting pain in the throat, so as to prevent swallowing without great inconvenience; when the glands of the neck are tumefied, tender to the touch, and impede deglutition; and when, in addition to these symptoms, we experience a sense of diffused coldness, increased towards night, and succeeded by heat. Absence of thirst is another indication of the propriety of administering *Pulsatilla*.

Belladonna is frequently in requisition in quinsy, and the symptoms which proclaim its use are the following:

Sensation as of a ligature round the throat, with external swelling; enlargement of the tonsils; bright redness of the entire throat, of the velum palati, and of the tonsils, with greater or less difficulty of swallowing; constriction of the voice; general and burning heat; much thirst.

In cases where the fever has been violent from the onset, we may commence our treatment by a dose or two of *Aconite*.

Belladonna, in one, two, or three doses, will subdue this disease; and, where any trace is left, *Mercury* will quickly succeed in the restoration to perfect health.

The following medicines are, equally with those mentioned above, entitled to important consideration: *Argentum*, *Barytes*, *Dulcamara*, *Nux Vomica*, *Rhus*, *Sabadilla*, and *Sepia*.

Croup.

This disease, with which most persons are conversant, is very important, rapid in its progress, and requires the most efficient means to avert its danger.

At its commencement, this affection is frequently deceptive, and assumes a catarrhal form. The patient complains of hoarseness, mental and bodily depression, slight febrile excitement, particularly towards evening, united with dry, short, hollow cough, with a slight sense of burning pain, and tickling in the wind-pipe. Soon, however, the sound of the cough, voice, and respiration assumes a special character, presenting nothing analogous to catarrh. It is veiled, sonorous, suspirious, whistling; sometimes grave and resonant, at other times shrill and squeaking; the respiration is more or less wheezing, and the cough excites pain, which the sufferer indicates by carrying his hand to his throat. Occasionally, however, the disease is sudden in its access, and not preceded by catarrhal symptoms. The cough is consequently hoarse, barking, hollow; or, on the other hand, acute, shrill, and clear; the voice is hoarse, rough, or wheezing; the face is red, mottled, swollen, and at times blue; a clammy sweat suffuses the body, the breathing is difficult, and the head is thrown back to facilitate respiration.

Aconite is the first remedy to be administered in this affection, in order to subdue inflammatory action.

After the exhibition of *Aconite*, one or two

doses according to circumstances, we must administer burnt *Sponge*. Should the symptoms refuse to yield to this remedy, we must afterwards have recourse to *Hepar Sulphuris*.

In ordinary cases, the sponge should be exhibited two or four hours after the aconite, and the repetition of the sponge and of the liver of sulphur should be alternated every two, four, or six hours. It is evident, however, that we must be guided by the progress of the disease.

Excoriation of the Nipples.

This affection attacks women recently delivered, and frequently becomes so painful as to prevent their giving the breast to the child.

The skin of the nipples is so delicate that the most trifling sores, upon the slightest contact, produce in many women almost insupportable suffering; and, discontinuing to suckle upon the affected side, the milk is suppressed, and inflammation of the mamma supervenes.

Women may prevent this affection by commencing, many weeks previously to their delivery, to wash the nipples every day with spirit distilled from corn.

They should continue a similar application

after their *accouchement*, so long as the nipples are without excoriation, taking the precaution to wash them with tepid water, every time the child is put to the breast.

Arnica is a valuable remedy where the excoriations are apparent. It is employed externally, in the dose of two or four drops of the tincture in fifty or eighty drops of water, taking care to wash the nipple every time the child is permitted to suck.

When the complaint is so severe as to present a certain degree of inveteracy, and especially where there is an evident psoric diathesis, we must have recourse to remedies administered internally. Of these *Sulphur* is one of the most important. Should it not entirely succeed, however, we shall, after its exhibition, derive most benefit from *Lycopodium*, *Calcarea Carbonica*, and *Graphites*.

If *Arnica* and *Sulphur* fail in immediate relief, we should consult a medical practitioner, since none else can discriminate the cases in which the other remedies may be applicable.

Excoriation of Children.

This complaint is sufficiently frequent in early infancy; and the non-observance of cleanliness,

moisture, and heat of weather, constitute in most instances the exciting cause. These excoriations are observable upon all those parts of the skin where folds are to be met with, and more particularly in places where there is much fat, as the neck, the hams, the thighs, the arms, and the genital organs. Proper attention to cleanliness will often succeed in preventing this disease, and the tepid bath is an excellent adjuvant. Nevertheless, we occasionally find it presenting itself in defiance of the most scrupulous attention, and this, too, without any assignable cause. Should the affection be trivial, cleanliness and the application of an innocent drying powder will quickly relieve it.

When the complaint is very severe, and causes much uneasiness to the patient, we must persevere in ablutions, and administer internally *Chamomilla*. In cases where there may have been an abuse of this remedy, *Pulsatilla* or *Ignatia* will be the appropriate agent.

When the disease assumes a more serious aspect, and the body of the child presents the appearance of a wound, *Mercury* will prove a specific.

Should the affection resist this means, we must attack it with antipsorics, and *Graphites*, *Sulphur*, and *Lycopodium* will then subdue the evil. The more serious species of excoriation have their

ultimate cause in a psoric tendency, and it is for this reason that antipsoric remedies will especially avail.

Catarrhal Fever.

This disease is frequently the sequel of a cold occasioned by a sudden change in the temperature; or from humidity in the atmosphere; and it is on this account that we so often find it prevailing in certain places and at certain seasons as an epidemic. If this affection be properly treated, it is of trivial moment, and becomes important only in consequence of the injudicious treatment to which so many persons are subjected.

The symptoms which characterize this condition are as follows:

Shiverings; transitory chills, shifting from one part to another, and alternating with a sensation of heat; pulse soft, and of moderate frequency; sense of heaviness, twitching or cracking in the limbs; the back and the sacrum feel as though they were broken; the head becomes confused, and is very weighty; pressive pain above the eyes; oppressive wakefulness, or disturbed slumber; diminution of taste and smell; the patient complains of the pit of the stomach, frequently feeling pressure coupled with anxiety; taste in the mouth

frequently bitter or mucous; appetite more or less deficient; the febrile symptoms especially augmented towards night; and the nose, throat, chest, and heart being considerably affected.

What will aid us in our diagnosis are the symptoms of catarrh: the nose, throat, and lungs secrete a vast quantity of mucus, feel very painful, and afterwards experience a sensation of heat. The mucus, at the commencement, is usually thin and white, but it subsequently becomes thick, viscid, and yellow, or even darker, and acquires a fetid odour.

The remedies most appropriate to combat this train of symptoms are Aconite, Nux Vomica, Chamomilla, Pulsatilla, Ignatia, and Belladonna. The constitution of the patient, the change in his moral tendencies, and that in the symptoms enumerated above, will guide us in our election.

Aconite is of service only in the commencement of the attack, and where the malady exhibits itself in an aggravated form: it acts in subduing arterial action.

Nux Vomica is almost invariably suited to patients whose temperament is quick and choleric; where the affection is accompanied by thirst and constipation; where the different parts of the body are under the influence of cold shivers, which, especially on moving about, appear to pass

over the bones, and alternate with heats. It is similarly eligible in cases where the symptoms are relieved by proximity to the fire; also where the patient endures a sense of scraping in the throat, hoarseness, and severe cough, more particularly in the morning.

Pulsatilla is suitable to persons of placid temperament, disposed to melancholy and to tears; to individuals of lymphatic habit; to women; and to patients who are but slightly inconvenienced by thirst. Also in cases where the secretions are too abundant, and become thick; where there is clamminess, as it were, of the mouth, loss of appetite, languor; and where, in conjunction with these, the other symptoms exist in a greater or less degree. This medicine, in certain instances, is appropriate after the exhibition of *Nux Vomica*, that is, where indicated by the change in the disease.

Chamomilla should be prescribed where there is impatience under suffering, irritability and extreme sensitiveness to pain. This remedy is very valuable for children.

Ignatia. It is frequently the moral symptoms which should guide the choice of the practitioner, and induce him to adopt *Ignatia*. This medicine is eminently suited to persons of gentle temperament, who pass rapidly from joy to grief, and

who are disposed to confine their griefs within their own breasts.

When to these moral tendencies are united symptoms of catarrhal fever, there is no doubt but that this remedy should be selected in preference to others.

Belladonna is suitable to children of sanguine and lymphatic temperaments, and affected with catarrhal fever; where the cough is evidenced chiefly at night; where the viscera are threatened with congestion; excessive irritability; somnolency during the day, with absence of sleep during the night; where the sleep is interrupted by sudden starts; delirium, more or less intense; prostration of strength; sadness; and where the patient screams out as though under the influence of fear.

Fatigue.

Although fatigue can scarcely be regarded as a disease, since rest, after a cold bath or ablution with cold water, will usually suffice to remedy this state, it is nevertheless desirable to point out the means which will quickly remove it, and avert the symptoms which occasionally follow in its train.

Arnica, in the dose of one or more globules, in-

ternally, and *Arnica* with water, as an external application to those parts of the body suffering from fatigue, such as the legs and feet, will speedily remove all inconvenience.

A decoction of hempseed, as a warm bath to the feet, will readily counteract any fatigue in those organs.

Milk Fever.

This fever exhibits itself in women two or three days subsequently to delivery. It is produced by a change effected in the organism secreting the milk. It supervenes with slight shivering, heat more or less ardent, anxiety, difficulty of breathing, twitching pains in the back and extending towards the breasts, headach, frequency of pulse, perspiration profuse and sour to the smell. This disease does not affect all women, although the majority are subject to its influence. They who are subject to febrile excitement are chiefly liable; also they who may be exposed to any accidental cause, such as a wound, contusion, protracted or severe labour, moral impressions, &c.

During the fever, the lochial discharge is diminished, but, if there be no complication, the patient is devoid of uneasiness in the lower part

of the abdomen. As soon as the secretion of milk is established, the fever subsides, and the lochia resume their full flow and primitive character.

Arnica is the remedy applicable to all cases where the labour has been protracted: where we find contusion or laceration, we may administer it externally in the form of lotion.

Aconite is indicated where the fever assumes an inflammatory character.

Bryonia is the remedy where the inflammatory action is less decisive, or where it has been controlled by *Aconite*; and where we find symptoms of rheumatic pain in the breast, externally and internally.

Belladonna should be given where the fever, more or less severe, is connected with an erysipelatous affection of the breast.

Emotions of the mind are frequently the exciting cause of milk fever.

Coffea is applicable in cases where the patient experiences a lively joy.

Chamomilla, preceded by *Aconite* when the fever is high, is an appropriate remedy, more especially where passion has been the excitant.

Opium is efficacious where the affection may be traced to fright or apprehension.

Aconite should be the remedial agent where

fright and passion have been the exciting causes of the mischief.

Concentrated grief, giving place to the disease, indicates the propriety of *Ignatia*.

We must be careful not to confound this affection with puerperal fever. The latter malady is of fearful importance, and demands the aid of a professional man. Thus, in all instances, where milk-fever presents alarming symptoms, and is conjoined with pain in the pectoral organs, abdomen, or head, we must instantly refer the case to a medical man.

Flatulence.

We do not in this place allude to chronic maladies which develop flatulence: our attention is directed merely to accidental cases, resulting from the use of flatulent vegetables; from having taken, in too large a quantity, cold liquids, or new and badly fermented beer; or from having partaken of indigestible food. Flatulence swells the abdomen, excites pain, impedes respiration, and causes general uneasiness.

China is often of great utility when colicky pains affect the lower intestines, and where the patient feels a sense of tightness; where the wind

is unable to escape ; and where, beneath the false ribs, there is tension and anxiety.

Nux Vomica will principally succeed among patients of lively temperament, and suffering from constipation.

Pulsatilla is suitable to women whose disposition is naturally mild and gentle ; and especially where flatulence follows the use of rich food.

Cocculus is applicable in cases where the patient experiences a sense of tightness and contraction in the lower region of the abdomen, and in the organs of reproduction ; uneasiness in the stomach ; where no relief is derived from the expulsion of wind ; partial swelling of the abdomen, with pressive, tearing, and burning pain ; also when the epigastrium becomes swollen ; when there exist gripings, twitchings, anxiety, and pressure under the cartilages of the ribs ; and when these symptoms are relieved by the removal of the offending matters.

Sprains.

By a sprain we understand a violent distention of the ligaments of a joint, originating in undue exertion, or in a fall. It exhibits itself by pain in the organ affected, and by weakness and swelling in the entire joint.

The first injunction is perfect rest; we may afterwards exhibit *Arnica* internally, and at the same time have recourse to friction with tincture of arnica to the injured part. This we may repeat three or four times a day.

When the arnica does not readily succeed, we may have recourse to *Rhus*, which may be administered in a similar manner, both externally and inwardly.

The dose of arnica and rhus is from one to three globules, in proportion to the exigency of the case. Four, or six, or ten drops of the tincture, in one hundred drops of water, may be employed for the friction.

Fright.

The lively and disagreeable impression derived from an impending danger imparts a greater or less degree of mischief to the nervous system, and the entire organism is more or less affected.

Fear may be succeeded by stupor, fainting, suspense of the action of the heart, numbness of the muscles, paralysis, apoplexy, and even by an alienation of the mental faculties. We may conceive that the effects of fright will widely vary according to diversity of constitution, individual

susceptibility, &c. For instance, persons who are delicate and of excitable temperament are very powerfully influenced by circumstances which would produce no effect upon the more robust.

Opium, taken immediately after the fright, is the best remedy for preventing any mischievous results, and more particularly so where we remark the following symptoms:

Tearing, frontal headach, pressing from within outwards; risings and acid vomitings; inclination to lie down; cold and sudden perspirations; sensation as of a weight in the lower part of the abdomen; anxiety; internal heat; stupor; sudden numbness of the body, with asthma; tremor of the limbs, with oppression and isolated startings of the body; convulsion in the limbs, with external coldness of the body; a species of apathy, with snoring, &c.

If some hours or more have elapsed since the accession of the fright, without having administered any medicine, we may give one drop of the tincture of *Sambucus*.

In cases where we find powerful action on the sanguineous system, we must have recourse to *Aconite*, subsequently to the administration of either *Opium* or *Sambucus*.

Boils.

A boil is a circumscribed, elevated, and pointed tumour, slightly indurated, of a deep red colour, and more or less painful, situated beneath the skin in the cellular tissue, and usually varying in size from that of a hazel-nut to a pigeon's egg, or even larger.

This tumour almost invariably terminates in suppuration: a kind of head is formed in the centre, after the removal of which there is a discharge of pus mixed with blood. The pains then subside, and all trace of the tumour gradually disappears. We may occasionally observe this affection healing without arriving at suppuration; but in such cases we almost invariably perceive an induration of the skin, and, notwithstanding there may be neither redness nor pain, the skin quickly swells afresh, and there is a relapse of genuine boil.

Arnica, repeated at intervals in small doses, will heal these furuncles, and frequently prevent their return.

Belladonna may advantageously supersede the *Arnica*, should the latter remedy fail in its results.

In cases where the boils originate in a psoric condition, *Arnica* will not suffice; we must therefore enforce the antipsoric treatment.

Swelling of the Glands.

Swelling of the glands situated beneath the chin, in the throat, or in the nape of the neck, results, among children of scrofulous diathesis, from the most trivial circumstance; and along its entire course the neck appears as though encircled with a chaplet.

Subjects disposed to this disease are all under the influence of the psoric virus, and it is only a prolonged medical treatment, suitable exercise, regimen, exercise in the open air, and habits of perfect cleanliness, which can remedy such constitutional tendency.

Dulcamara, *Mercury*, and *Belladonna*, almost invariably succeed in reducing the glands to their natural size. An experienced practitioner should always be consulted in pursuing a radical course of treatment.

Hemorrhage.

Hemorrhage is the name assigned to a loss of blood which may be sufficient in quantity to threaten the life of the patient. The discharge takes place most frequently from the mouth, and, among women, from the uterus. Under the former circumstance the blood may proceed from the lungs

or from the stomach. When the hemorrhage is from the lungs, it is heralded by a sense of heaviness and constriction in the chest, asthma, palpitation of the heart, anxiety, ebullition of blood, cough, and a sweetish taste in the mouth.

The spitting of blood occurs after a slight attack of cough, and sometimes after a tickling sensation in the upper portion of the chest, or in the throat, which inclines the patient to cough.

The most suitable remedy, and one to be administered as quickly as possible, is *Aconite*, which may be repeated every two or three hours, should circumstances threaten a relapse.

China is applicable after the exhibition of *Aconite*, when the sanguineous excitement is calmed; when the patient feels shivering or transitory heats, fatigue, or weakness; when he is inclined to lie down; when the limbs are tremulous, brief sweats exhibit themselves, and the party continues to expectorate blood, more or less, every day: it is in the intervals between the hemorrhages that we should administer this remedy.

Pulsatilla is appropriate where the patient feels worse at night than by day; when he complains of shiverings, with sense of general debility; of uneasiness in the lower portion of the chest, shootings in the chest, and difficulty of respiration; when the blood expectorated is deeply coloured, and in

clots; when the feet are cold, and the patient is overwhelmed with grief, and inclined to tears.

Pulsatilla is very efficient among women presenting the above symptoms, and who labour under suppression of the menses.

A medical man should always be consulted in cases of hemoptysis, and we merely indicate the first steps to be taken previously to his arrival. The remedies next in importance to those we have pointed out are the following, while the study of the *Materia Medica* and the diversity of cases will lead the physician to the choice of the most eligible remedy:

Acid. Sulph., Arnica, Belladonna, Crocus, Digitalis, Drosera, Kali, Ledum, Lycopodium, Millefolium, Rhus, Arsenic, Ferrum, Carbo Veg., Phosphorus, Ammon. Carb., Calcareo, Magnes. Carb., Plumbum, Acid. Nitric.

When the blood proceeds from the stomach the premonitory symptoms are general uneasiness; heaviness in the limbs; confusion in the head; anxious sense of pressure in the epigastrium; swelling and fulness of the abdomen; costiveness; disposition to vomit, with mawkish and putrid taste in the mouth. The blood which is discharged by the vomiting is of a deep red or blackish colour, and is frequently grumous.

Nux Vomica should be given where the above-

mentioned symptoms are observable; we must be careful to prescribe a very weak dose, so that we may repeat it every three or four hours, according to the progress of the disease, while awaiting the arrival of the medical man.

The medicaments appropriate to this disease, and which the practitioner should enquire into, are—

Aconite, Arsenic, Hyoscyamus, Ipecacuanha, Veratrum, China, Belladonna, Stannum, Arnica, Millefolium, Cantharides, Mezereon, Cicutā Viridis, Sulphur, Lycopodium, Zincum, Calcareā, and Carbo Vegetabilis.

Uterine hemorrhages are the most frequent, and are particularly dangerous when they occur during pregnancy.

The circumstance of pregnancy renders it essential to enjoin perfect repose.

Ipecacuanha is the first remedy to be adopted, provided the hemorrhage be not associated with uterine contractions, or pains similar to those of labour, and where the loss of blood is abundant, continual, without gushings, and accompanied by colic and general prostration.

Chamomilla is suitable where the symptoms of the disease consist of loss of blood, with pains resembling those of labour. This remedy is

especially indicated where anger has been the exciting cause of the disease.

China is an appropriate remedy where the patient feels uterine contraction, with pains resembling those of childbirth, and extending towards the rectum; if the hemorrhage come on in paroxysms, increasing and diminishing with the pains; if the patient feel colic, with painful tension of the lower part of the abdomen; if the patient be weakened for the moment; be disposed to fall, and really do fall, into a state of lipothymy; if the skin be cold, and she have a pressing inclination to make water.

Hyoscyamus is eligible where the uterine discharge is connected with general excitement of the nervous system; tremor of the whole body; pains resembling those of childbirth; giddiness; agitated slumber; great inquietude; excessive vivacity; convulsive movements; general heat; swelling of the veins; extreme violence of pulse; alteration of feature, &c.

Arnica should be prescribed where the symptoms are referrible to external violence, undue exertion, &c.

Crocus is applicable where the blood which escapes is black, clotted, and viscid, and where the patient feels cutting pains in the lower part of the abdomen, which extend through to the back.

Sabina is a specific where the blood is of a bright red colour, and is expelled by the contraction of the labour-pains.

Belladonna is very salutary where the hemorrhage is accompanied by pressure from above downwards, and from behind forwards, in the organs of generation; when the blood is bright red; where there is agitation, with hard, full, and frequent pulse.

We may still further consult the following medicaments: *Bryonia*, *Platina*, *Stramonium*, *Ignatia*, *Ferrum*, *Pulsatilla*, *Carbo Animalis*, *Secale Cornutum*, and *Sepia*.

Hemorrhoids.

This very frequent affection has its first cause in a psoric state of the economy, which by development effects a change in the circulation of the fluids in the vessels of the lower region of the abdomen, after derangement, more or less evident, in the functions of the abdominal viscera. It principally affects men of middle age, and women after the suppression in their monthly periods: it has been observable even in infancy.

This disease is characterized by congestion in the head; vertigo; congestion in the chest; palpi-

tation of the heart ; pains in the back ; tenesmus ; itching at the anus and in the rectum ; obstinate costiveness ; colics ; and, lastly, by an excrescence more or less developed around the anus. It reappears at fixed or indeterminate periods, and usually terminates in a discharge of blood or mucus more or less abundant. At times, however, this excretion does not take place, and the hemorrhoids or piles are termed *dry*.

This disease is frequently rendered incurable from the injudicious treatment to which the patient has had recourse, or from persevering in a regimen altogether at variance with the malady.

The sufferer should consult a competent medical authority with reference to an antipsoric plan of treatment to relieve, if not too late, so painful and distressing a disease.

To re-establish his health, the patient must submit to a very moderate dietary : he must take plenty of exercise in the open air ; avoid colds, particularly in the feet ; he must abstain, as much as possible, from wine, coffee, tea, strong malt-liquor, lively emotions, late hours, and too assiduous application.

Nux Vomica is the most suitable application where the occasional causes of the disease are the use of stimulating beverages, such as wine, brandy, &c. ; late hours, toils of the study, cos-

tiveness, worms, pregnancy, swelling of the abdominal viscera, excrescences of considerable size, and which are the seat of burning and shooting pains; sense of strangulation in the rectum; contraction of this viscus, united with dull and shooting pains felt in paroxysms in the loins and in the bones of the ham; cracking pains in the loins upon the slightest movement, inducing the patient to utter cries, and preventing his holding himself erect, or walking in any but a bent position; flow of clear fluid blood after the alvine evacuations, or under other circumstances, but always accompanied by a desire to relieve the bowels.

Arsenic is very efficacious where the hemorrhoids present, as a leading symptom, a violent burning in the rectum, the hemorrhoidal tumours, and in the surrounding parts. The evacuations nearly aqueous, and a rapid loss of power, united with the above symptoms, still indicate the same remedy.

Capsicum frequently succeeds in the removal of the distressing pruritus of the hemorrhoidal tumours; when the patient experiences a burning sensation, more or less violent, and where there is diarrhœal discharge.

Belladonna is valuable in moist piles, accompanied by pain in the loins, which are nearly

insufferable, and with a sensation as though that part were broken asunder.

Sulphur is especially adapted to the following train of symptoms :

Continual desire to go to stool, even after diarrhœal or sanguineous motions ; darting pain of excoriation in the rectum, and neighbouring parts ; itching ; burning ; moisture of the excrescences ; occasionally a pressive fulness in the rectum, and frequently even the prolapsus of the part ; violent shooting pains in the loins, with sense of stiffness and tension, as though the skin and the muscles were contracted.

We may again, according to the circumstances of the case, derive the best results from the following remedies :

Carbo Vegetabilis, Causticum, Calcarea, Graphites, Kali, Lycopodium, Pulsatilla, Phosphorus, Acid. Nit., Colocynthis, Mur. Acid.

Catarrhal Inflammation of the Throat.

A scrofulous diathesis is one of the causes especially predisposing to this malady. It consists in an inflammatory affection of the mucous membrane which lines the throat, and also of the ganglions of the part. It exhibits itself chiefly

after exposure to colds, errors of diet, &c. It is characterized by redness, swelling of the mucous membrane, hoarseness, difficult and painful deglutition, and by a tickling and irritation in the throat. This disease very frequently affects the individual, by reason of the predisposition indicated above.

Chamomilla is suitable where there is dryness of the throat, with thirst; disagreeable and uneasy sensation in swallowing, or in bending the neck; feeling in the throat as of a foreign body, from which the patient desires to be relieved by spitting; swelling of the glands of the lower jaw, with pulsative pain; fever towards evening; alternations of cold and heat, with ill-humour.

Mercury is appropriate where an accumulation of saliva induces the patient frequently to spit; where a sense of burning in the throat renders deglutition difficult; the throat appears too much contracted; a smarting pain is experienced, extending to the ear; the almonds of the ears are swollen; a darting pain in swallowing; unpleasant taste in the mouth; swelling in the posterior part of the tongue and of the gums; access of catarrhal fever in the evening, with alternations of cold and heat.

Pulsatilla is an eligible remedy where the following symptoms present themselves:

Sense of excoriation, incisive pain, and dryness of the throat; swelling of the glands; shivering, increased towards night, and succeeded by heat; absence of thirst.

Belladonna is suitable where, in drinking, there remains in the throat a cramp-like feeling of strangulation, and where the throat is swollen externally.

Coffea may be administered under the following circumstances:

Continual pain extending from one side of the palate to the pharynx, and which is increased by deglutition; swelling of the superior portion of the *velum palati* and of the uvula, which is elongated; sense of mucous accumulation in the affected part, obliging the patient always to swallow; sensation of dryness and heat in the throat; inclination to cough; cold in the head; aggravation of symptoms in the open air; sleeplessness; heat; sensitiveness; melancholy temperament.

Cerebral Inflammation.

This disease is of frequent occurrence among children, and at an early period of life it is of fearful import: at all ages it is highly dangerous.

Having attained a certain height, this affection

yields with the greatest difficulty even to the most skilful treatment; and, for this reason, it is of the most signal importance to be intimately conversant with the premonitory symptoms. These are chiefly the effects of congestion in the head, producing pain and heaviness in that organ; change of disposition; fits of passion; anxiety; pains in the ears; intolerance of light; disposition to costiveness; frequent nausea, succeeded at times by vomiting; coma; inability to fall asleep, or fright during sleep.

The earlier symptoms are the following:

Heat and heaviness in the head; face very red and swollen; eyes suffused and brilliant; excess of irritability and sensitiveness; giddiness; state of coma; delirium; divers indications of spasm; grinding of the teeth; inclination to thrust the fingers into the nostrils, and to bury the head in the bolster; to seize with eagerness every surrounding object; visible throbbings in the veins of the neck; absence of thirst with great heat; forcing down of the lower part of the abdomen; constipation; occasionally, slight diarrhœa.

This disease demands the most prompt assistance.

Aconite is appropriate at the onset of the attack, in order to modify the sanguineous congestion in the head; we must not, however, dwell too long

upon the application of this remedy, which, in itself alone, does not possess the power of arresting the malady.

Belladonna is the remedy which renders the most essential service after the exhibition of *Aconite*, provided the latter medicine have been deemed necessary: it is eligible where the symptoms already enumerated present themselves.

Hyoscyamus is appropriate when the patient lies in bed without cognizance, with the eyes closed; the mouth opening with difficulty; the tongue loaded with white and frothy mucus; raves upon his own private affairs; sinks down without cognizance; stutters inarticulately; smiles; or is at times perfectly quiescent; moves his hands as though to withdraw the coverlid; frequent subsultus; dilatation of the pupil; fixedness of vision; redness of the face; skin dry and disposed to crack; respiration anxious and weak; full pulse, and tension of the abdomen.

Opium should be prescribed where the patient is in a soporific state; where the somnolency is accompanied by snoring; the eyes half open; the patient utters no complaint; is in need of nothing; and has frequent vomiting.

Stramonium is applicable where the patient, slumbering naturally, has starting in the limbs;

snorings; his movements irregular; and, upon waking, he appears without cognizance; sometimes fixing his attention; sometimes withdrawing himself slowly, and with apprehension; sometimes retreating with exclamations of dread; the above symptoms associated with ardent fever, redness of the face, and moisture of the skin.

Arnica, Sulphur, Digitalis, China, and Bryonia, may, under certain circumstances, be prescribed with advantage: it is for the practitioner to distinguish the peculiarities which should indicate their employ.

Inflammation of the Eyes.

In children recently born, a too lively impression of light occasions an inflammation of the eyes, characterized by a secretion, more or less abundant and viscid in character, from the mucous membrane of the eye.

Aconite, aided by the observance of cleanliness, will quickly succeed in the removal of this malady. In cases where the intolerance of light is excessive, where the conjunctiva is highly inflamed and turgid, *Belladonna* may be administered with great advantage after the *Aconite*.

Sulphur, succeeding to *Aconite* and *Belladonna*, will complete the treatment, should any trace of the disease remain.

Calcareo Carbonica may succeed to the exhibition of *Sulphur*, should the disease evince a tendency to the chronic state.

Inflammation of the eye, originating in the introduction of a foreign body, requires for its treatment the careful removal of the substance, to bathe the eye with pure water, and to administer *Aconite*, which may be repeated according to circumstances. *Sulphur* may follow this remedy, if the affection be not effectually removed by the *Aconite*.

Arnica is very valuable when the eye has received any severe contusion: if it do not effectually prevent the development of inflammation, we may have recourse to *Aconite*.

Orgelet, or *Stye*, is an inflammatory affection of the glands of the eyelid. It gives rise to a yellowish tumour, of the size of a grain of barley, and frequently causes pain in the surrounding parts. In most instances we gain prompt relief through the agency of *Pulsatilla*, administered internally, as also in the form of lotion.

When, as a sequel to such repeated inflammations, there may remain a slight induration of the lid from a thickening of a small gland, with

disposition to renewed inflammation—notwithstanding such tendency is not fully developed, but assumes a subacute form, so that the edges of the eyelids are always agglutinated—we may administer *Staphysagria*.

Catarrhal Ophthalmia

Proclaims itself by redness of the eyelids, and sometimes of the conjunctiva, more especially at the angles of the eyes; there is also a sense of burning and compression, as from the presence of a grain of sand, dread of light, and discharge of tears: to these symptoms are frequently united catarrh, with or without fever, and a dry cough, which is more or less troublesome.

Aconite is the first remedy by which to arrest the inflammatory progress, and may be followed up by *Nux Vomica*, provided the affection be not too severe.

Belladonna is appropriate after the *Aconite*, and where the inflammation of the mucous membrane is intense, the vessels highly injected, and the eyes very sensitive to light; where we find a violent cold in the head, excoriating the nostrils and alæ of the nose, the lips, and their commissure; small inflamed pustules; and cough returning

at intervals, with a sense of suffocation and wheezing.

Euphrasia is a preferable remedy where the redness of the lids,—produced by violent cold, with pain in the head, and exacerbation of fever towards night,—extends itself over the whites of the eyes: this remedy is especially indicated where the secretion of tears is abundant.

Ophthalmia presents an infinite number of variations which we cannot discuss in this brief treatise: we confine ourselves therefore to those of most frequent occurrence.

The following are the more important remedies to be administered in affections of the eye:

Acid. Nitric., Acid. Sulph., Arsenic, Belladonna, Calcareæ, Causticum, Chamomilla, China, Colocynthis, Conium, Digitalis, Clematis, Euphrasia, Graphites, Hepar. Sulph., Mercury, Nux Vomica, Petroleum, Psoricum, Pulsatilla, Rhus, Sepia, Spigelia, Sulphur.

Inflammation of the Lungs.

Inflammation of the lungs evinces itself most frequently without any premonitory symptoms: we observe violent shiverings, succeeded by in-

tense heat; and it is in most instances as a sequel to cold that this affection presents itself.

The principal characteristic symptoms are pressive and darting pains in the chest, considerably aggravated by inspiration; great oppression; shortness of breath; anxiety; cough, attended with much pain, dry at first, and succeeded by sputa tinged with blood. On percussion, the pain in the chest is not sensibly increased, but the parts to which percussion has been applied, and corresponding to the affected portion of the lungs, present a sound more or less dull, whilst the healthy parts are clear and sonorous.

The stethoscope, or even the ear applied to the chest, enables us to detect a peculiarity of sound which indicates perfectly the part affected; and where the pneumonia has attained a high degree of intensity, the lung ceases to be permeable to the air, and the respiratory sound is no longer perceptible. The pulse is full, strong, and frequent, oftentimes with palpitation of the heart; the skin is hot, dry, and burning; the face is red and swollen; the urine is high coloured; excessive thirst, and, in most instances, constipation more or less obstinate.

As soon as the above details are observable, we should administer *Aconite*, two or three globules to the dose; and if there be no favorable change

in the general symptoms, or in the state of the circulation, we may repeat the medicine. We need be under no apprehension of exhibiting two, three, or four doses of aconite, during the first twenty-four hours.

This remedy alone will frequently avail in arresting the progress of the mischief. *Bryonia*, however, is oftentimes indispensable after the administration of *Aconite*; and, according to circumstances, we may give a dose every four, six, or ten hours. This remedy is especially indicated where there are darting pains in the chest; where the sputa, tinged with blood, present a rusty appearance; and where we find rheumatic pains in the back, with oppression.

The practitioner will readily perceive that *Rhus* is especially suitable in pneumonia, after the use of aconite and bryonia, and where a typhoid condition is apparent.

Squilla will be serviceable where the *Aconite* has removed the earlier symptoms, and where there remains a profuse mucal expectoration; also where the patient complains of weight beneath the sternum, with lancinations in the right side of the chest, deeply towards the back; respiration very short, quick, and anxious; the patient lies in bed with the shoulders much elevated; the pulse slightly

defined; evening chills without being succeeded by heat, together with absence of sleep.

This remedy, then, is adapted to circumstances less decidedly inflammatory than where *Bryonia* would be appropriate.

In an elementary work like the present, we have no room to indicate the less frequent varieties to be met with in pneumonia: we shall content ourselves with pointing out the remedies to which the practitioner should direct his particular attention, as being at times of essential importance. These are:

Arnica, Arsenic, Belladonna, Cannabis, China, Pulsatilla, Sulphur, Acid. Phosphoric., Nux Vomica, Hyoscyamus, Stramonium, Veratrum, Opium, Tartar. Emetic., Dulcamara.

Inflammation of the Breast in Nurses.

This disease originates in a predisposition peculiar to suckling; in errors of regimen, colds, lively emotions, fright, passion, mechanical lesion, such as a blow, and above all in weaning the infant too suddenly. In such cases, it frequently happens that the milk coagulates in the lacteal vessels; tumefaction more or less considerable occurs; the skin becomes tense, indurations are

detected in different places; an erysipelatous inflammation supervenes, almost invariably terminating in suppuration, which discharges from without, and causes most agonizing suffering to the patient. We frequently observe the establishment of renewed indurations, which terminate successively in a similar manner. This disease may thus continue several weeks, or even months, and terminate at length in deeply-seated cicatrices, and even in indurated tumours.

Belladonna should be administered immediately upon the skin becoming erysipelatous, and attended with gradual tumefaction of the mamma.

Bryonia is very suitable where the secretion of milk is suspended,—where there is tumefaction of the lacteal vessels,—where the bosom is obstructed,—and where the skin can scarcely be considered as erysipelatous. The medicine may be repeated according to the result.

If the inflammation be extensive and erysipelatous, with considerable swelling, and suppuration is commencing, we may give phosphorus; and finally, in order to facilitate the opening of the abscess, now fully ripe, we may give one or more doses of *hepar sulphuris*.

If the disease have already existed some time, if it have been injudiciously treated, and suppuration goes on without destroying the inflammation

of the breast, we must have recourse to *Silicea*, which, in one or more doses, will quickly restore the patient to health.

Vigilance.

Vigilance in adults, when unaccompanied by any other indication of perverted health, is in most instances the result of undue excitement of the nervous system, originating in intensity of thought, powerful emotion, the abuse of stimulating drinks, coffee, &c.

Our treatment must commence by causing the patient to abstain as much as possible from the exciting causes, and a dose of *Nux Vomica* may be administered in the course of the evening.

Pulsatilla is suitable where the wakefulness proceeds from overloading the stomach, giving rise to flatulence and constipation.

Wakefulness in young children frequently alarms their parents, and results, as a matter of course, from some derangement in their health. It frequently originates in errors of diet, and in such cases it is accompanied by swelling of the lower part of the abdomen, flatulence, green evacuations, &c. *Chamomilla* will be attended with perfect success in retrieving the mischief, and restoring

the patient to perfect health. Should this remedy, however, have been abused, we must have recourse to *Coffea*.

The vigilance which results from weaning will yield to *Belladonna*, if in the course of a few days the evil do not cure itself.

Wakefulness, in connexion with continued crying, during which the infant contracts the legs upon the abdomen, tosses about incessantly, and suffers severe griping pains, with diarrhœa, has its remedy in *Cina*.

Vigilance, resulting from a complicated state of disease, requires the treatment of the affection of which the wakefulness is merely symptomatic, and such prescription would be irrelevant here.

Irritability of the Skin.

There are individuals whose skin is so excessively irritable, as to become inflamed and suppurate under the influence of the most trivial causes. This diseased condition depends upon some peculiar predisposition, and rarely gives way except to an antipsoric treatment long persisted in. *Chamomilla*, *Hepar Sulphuris*, and *Petroleum*, may be successfully opposed to this condition of the skin, and may be varied according to the exigencies of the case.

Intoxication.

Vomiting, being an excellent means of preventing the mischievous results of inebriety, may be facilitated by drinking warm water, or a strong decoction of *Coffea*. Very frequently, however, *Nux Vomica* will succeed in dispersing the symptoms indicating such a state of things, and prevent the dependent consequences which are more or less distressing.

Jaundice.

This affection results from innumerable causes, and presents itself under many different aspects.

One of the causes, most frequent in occurrence, is passion: the skin assumes a yellowish appearance, as also the whites of the eyes, and the alæ of the nose; the urine becomes brown and thick; the abdomen contracted; the evacuations white, grey, and slimy; the appetite fails, and the mouth acquires a bitter taste. We commence our treatment by *Chamomilla*, and after its use we administer *Nux Vomica* and *Bryonia*.

In cases where we meet with fever, after the

exhibition of *Chamomilla*, we shall do well to administer one or two doses of *Aconite*.

If the jaundice have originated in the abuse of *Chamomilla*, the symptoms must be combated by *Pulsatilla* and *Ignatia*, and the treatment concluded by *China*.

This disease among newly-born infants very frequently originates in cold, to which children, during the first few days after birth, are particularly liable; it may also proceed from the abuse of purgatives administered for the purpose of discharging the meconium.

The treatment under such circumstances is similar to that indicated above. *Chamomilla* and *Nux* readily mastering the disease. A warm bath every day, for a few minutes, will prove of admirable assistance in promoting the health of the child.

The remaining remedies, which prove of infinite service in jaundice, and which should be scrupulously investigated by professional men, are:

Arsenic, Belladonna, Mercury, Sulphur, Digitalis, Calcareo, Carbo Vegetabilis.

Fainting Fits.

We thus denominate a train of symptoms presenting numberless varieties, as well in the symptoms themselves, as in the importance of their results.

The following phenomena are observable:-

Debility, giddiness, and confusion in the head; twitching in the ears; dimness, obscurity of vision, or the sensation as of a veil before the eyes; the patient trembles, becomes pale, and experiences anxiety; a cold sweat displays itself upon the forehead and neck; the eyes lose their lustre, and are surrounded by a blue circle; the animal powers diminish, and the patient at length loses all cognizance and sense of feeling. On the other hand, there may remain merely a bewilderment of sense, confusion, and indistinct noise in the ears; the pulse weakens, becomes slow, small, and at times nearly imperceptible; and the respiration corresponds in all particulars with the pulse.

The more serious faintings, which very nearly resemble dissolution, consist in the complete suspension of the corporeal functions. Not only is there no cognizance, sense of feeling, or voluntary motion, but the pulse and the breathing, together with the excretions and secretions, are altogether suspended.

The irritability and sensitiveness, however, are not so far gone but that we may have recourse to galvanism, applied immediately to the denuded nerves and muscles, in order to re-establish the vital functions. This state of things is to be distinguished from death in reality, only by the

repeated application of means the most energetic, and by the absence of incipient decomposition.

The treatment of persons fallen into a swoon consists in the application of external excitants, such as friction to the soles of the feet, tickling the nose with a feather, and sprinkling the face with cold water.

A remedy which in most instances will readily succeed, consists in placing the patient horizontally, with the head rather lower than the body and the limbs. It is premised, however, that this inclination must be very trivial, and never put in force where there is any determination of blood to the head. It remains to observe, that these swoonings are of rare occurrence, and that a few moments will generally suffice to restore the patient.

The medicines, in such cases, are usually administered through the medium of olfaction, but we may also place the remedy upon the tongue. One of the best remedial agents is camphorated spirits of wine, which may be placed beneath the nose of the patient, and with which the forehead and temples may be slightly rubbed.

Opium is an appropriate agent where the swooning has been the result of fright.

Moschus is suitable in fainting fits, to which men who are hypochondriacal, delicate, and irritable,

and women of hysterical habit, are very liable; also young girls of a chlorotic tendency.

China is the remedy suitable to fainting-fits which result from a considerable loss of the juices, such as prolonged nursing, repeated blood-lettings, profuse hemorrhage, frequent pollutions, and long-continued diarrhœa.

Lypothymy, frequent in occurrence, and supervening without any appreciable cause, is referrible to some chronic ailment which must be confided to the care of the medical practitioner.

Fetid Breath.

As feter of the breath originates in most instances from want of paying proper attention to the teeth, the observance of cleanliness will remove the annoyance.

In cases where the odour proceeds from the stomach, or from the lungs, we must oppose it by the following medicines, provided the effluvia be perceived immediately on rising: *Carbo Animalis*, *Agaricus*, and *Iodium*.

Ammonium is the remedy where the patient himself remarks the feter of the breath.

Barytes, where the breath is very offensive; but the patient himself unconscious of the annoyance.

Graphites, where the odour resembles that of the urine, and proceeds from the nose.

Sulphur, if the odour become more offensive after a repast.

The fetid effluvia from the mouth are often the result of a species of putrid inflammation of its interior, in which the gums, tongue, and palate are affected with painful swelling, redness, burning heat, and emanation of putrescent smell with which the saliva is charged: the gums become soft, recede from the teeth, and are covered with small ulcers, the source of the fœtor.

Mercury triumphs readily over this species of disease, by restoring to a healthy condition the parietes of the mouth.

Aurum frequently removes fœtor of breath, particularly among young girls about the period of puberty.

Pains in the Loins.

This affection, often extremely violent, has its seat in the vicinity of the lumbar vertebræ, either in the lumbar region itself or in the sacrum; it possesses the same characteristics in either of these different parts, and is not increased by pressure.

At times the affection is confined to one side; flexure of the back and of the thigh is exceedingly painful; the sufferer is compelled to lie upon the back, or upon the affected side; he can neither turn, nor raise his thighs, nor separate them; when out of bed, he is obliged to walk constantly in a curved position; he frequently halts in his gait, and is unable to lift any burden; he complains of a tearing pain affecting the loins when he is desirous of standing upright, and which renders him nearly incapable of assuming that position. There are no external indications. This pain proceeds in most instances from a cold, excess in diet or drinking, and is frequently accompanied by a hemorrhoidal affection.

Nux Vomica is especially applicable where the symptoms have resulted from the causes above detailed; and where we find constipation with disposition to hemorrhoids. Where the disease is referrible to some severe chronic affection, it requires an antipsoric plan of treatment.

Arnica is an appropriate remedy where the disease has presented itself after raising a heavy weight, after a fall or contusion, and where the patient feels the pain so violent as scarcely to admit of his coughing, sneezing, or walking.

Should the *Arnica* prove insufficient, *Rhus* will succeed in perfecting the cure.

*Acute Inflammation of the Lumbar and Psoas
Muscles (Lumbago Inflammatoria).*

This may be distinguished by the following characteristics :

Pain, fixed, violent, drawing, tearing, tensive, and inflammatory, in the lumbar region, and extending itself at times to the back and shoulders, and even to the bladder and the thighs, without following the course of the nerves. It is frequently associated with colic ; a sensation at the anus, as though inclined to go to the closet ; cramps in the bladder ; dysuria, and strangury ; afterwards a sensation of debility, numbness, and torpor of the thigh on the affected side ; the patient cannot, except with difficulty, extend or flex the limb ; and, the pain increasing, he can scarcely lie upon his back in consequence of his suffering. Sometimes there is hardness, with evident swelling ; more or less febrile excitement accompanies the symptoms, which commence in a severe fit of shivering, followed by violent heat ; the skin is burning to the touch ; there is lively thirst ; and the pulse is hard, full, and incompressible.

The treatment should commence by mastering the general symptoms, and by exhibiting *Aconite*,

which may be repeated several times according to the exigency of the case.

Bryonia and *Nux* are serviceable where the pains are more especially experienced during motion.

Rhus and *Pulsatilla* may be employed where the patient suffers principally during quiescence.

Cantharides, where we find spasmodic symptoms in the urinary organs, such as difficulty in passing the water, or inability to micturite.

Mercury and *Chamomilla*, where the pains are almost insufferable during the night.

Staphysagria will prove efficient where the pains are pulsative in their character, and where there is a tendency to suppuration.

In such cases the following medicines will frequently be of service, more particularly where the pains are drawing, tearing, and inflammatory in their character :

Arnica, *Chamomilla*, *China*, *Mercurius*, *Hepar Sulphuris*, *Aurum*, *Assafoetida*, *Belladonna*.

The above remedies, as well as *Nux* and *Rhus*, are valuable both in acute and chronic cases.

Digitalis, *Colocynth*, *Argentum*, and *Plumbum* are more particularly appropriate to cases which are chronic in their character.

Cholera.

Cholera in this climate is most frequently observed during the prevalence of the summer heats, and assumes an epidemic form; it sometimes offers a complication of symptoms, and presents itself under a spasmodic or bilious type.

The predisposing and occasional causes are, in most instances, an elevated grade of temperature long continued; a sudden decline in the thermometer during a very hot day; peculiar irritability of constitution; a state of hysteria and hypochondriasis; the period of dentition; violent emotion, fear, grief, vexation; undigested aliments; acid, unripe fruit; cold drinks; coldness of the feet, and of the lower part of the abdomen; suppression of the natural discharges, such as the menses, perspiration, &c.; or even the exhibition of violent remedies.

Cholera not infrequently supervenes unattended by any precursory symptoms; or before the development of symptoms characterizing the affection, we may observe various phenomena, such as an increase of the bilious secretion; corporeal inactivity; general uneasiness; heaviness; jaundiced complexion; the tongue clothed with yellow mucus, and the root especially loaded and yellow; a bitterness of taste in the mouth; aversion for food;

bitter risings ; pressure and cramp-like contraction of the stomach ; sensation of fulness at the pit of the stomach ; anxiety in the epigastric region ; flatulence ; distaste ; swelling of the abdomen ; a rumbling and colic in the intestinal canal ; the urine deposits a reddish sediment, smells offensively, and pain is caused by the act of micturition.

When these symptoms are not relieved by appropriate remedies, or when the disease is sudden in its access, we remark the under-written phenomena :

The patient vomits his food, the ejection of which is succeeded by the more or less abundant discharge of a mucous, watery, and bilious fluid ; the colour of the vomited matter is yellow, green, brown, or blackish, and frequently very offensive. At the same time violent diarrhœa is established : at first the secretions are fecal, watery, and bilious, frequently associated with violent colicky pains, burning and acute in character, and occupying chiefly the region of the umbilicus.

Should the malady continue its progress, other symptoms present themselves : these are, fulness in the epigastrium ; quick and anxious respiration ; violent cramps in the stomach, with a spasmodic pulse, at times nearly imperceptible.

These symptoms frequently acquire extreme

severity; the strength and the pulse sink rapidly; spasmodic symptoms present themselves in different organs; superadded to which are painful cramps in the limbs; spasms in the bladder; cold perspiration; syncope, with hypocratic expression of countenance.

The preliminary symptoms of cholera are speedily removed by the administration of *Chamomilla*.

Ipecacuanha is appropriate where the vomitings and diarrhœa are already established. When the first doses of this medicine do not effectively meliorate the disease, or where it has acquired a high degree of development, we must unhesitatingly have recourse to *Veratrum*, which will act altogether as a specific.

Arsenic is required, when, to the characteristic symptoms of cholera are united a rapid declension of animal power; much anxiety; insatiable thirst; almost constant discharge from the bowels; the matters tinged with blood; vomitings; excruciating pain in the bowels; pulse scarcely perceptible; hypocratic expression.

Chamomilla is suitable to that form of cholera which is excited by a fit of passion.

Pulsatilla may be employed, where the evacuations are especially mucous, and will succeed perfectly well in subduing the affection.

The repetition of the medicine is of the utmost importance in this disease, since we must bear in remembrance, that frequently the virtues of the medicament, and vital reaction, are promptly extinguished by the morbid virulence. We should therefore repeat the remedy in accordance with the urgency of the case. In furtherance of this view, we should dissolve from one to six globules in twelve spoonfuls of water, and administer a spoonful of the mixture every quarter of an hour, every half-hour, or every hour, in proportion to the severity of the symptoms. In some instances of extreme character, it will be necessary to repeat the medicine every ten minutes. It is not the dilution of the remedy which is important, but the frequency of its repetition.

Medical practitioners will do well to direct their attention to the following medicines, in the event of those indicated above not corresponding with the cholera falling under their observation, such, for instance, as Asiatic Cholera.

China, Colocynthis, Dulcamara, Camphora, Cantharides, Carbo Vegetabilis, Cicuta, Belladonna, Cuprum, Lauro-Cerasus, Nux Vomica, Secale Cornutum.

Food in cases of cholera should be interdicted, since all aliments will be rejected by the stomach. The drinks should be iced, and administered in

small quantities at a time. If ice cannot be procured, the drinks should be given as cool as possible.

The reaction of the skin may be promoted by dry friction of the limbs. The temperature of the apartment should be moderate, as also that of the bed; and it must be remembered that all external applications of a medical character are perfectly futile. The internal exhibition of suitable remedies will promptly remove the evil.

Measles.

Measles is an eruptive disease attacking individuals of every age, but more especially those of tender years. It has its origin in some peculiar condition of the atmosphere, of the nature of which we are totally ignorant, and reigns chiefly in an epidemic form: cases purely isolated have rarely come under observation.

This disease is in many instances slight, and presents but little to excite apprehension; at other times it may be characterized by symptoms of extreme danger, and it leaves most formidable traces of its virulence. We may rest satisfied, however, that, under the influence of a judicious homœopathic treatment, this affection may be promptly and radically cured. It is almost un-

exampled to remark an unfavorable termination, except where the patient is found under circumstances altogether adverse, such as shattered in constitution, in abject misery, in a state of complicated psora, or where there is an utter deficiency of hygeinic observance.

Measles is invariably preceded by a catarrhal and febrile state, generally supervening three or four days previously to the appearance of the malady. The patient feels languid, complains of uneasiness, debility, of chills alternating with heats, and frequency of pulse; there is slight cough, with hoarseness and difficulty of respiration, accompanied with coryza; the eyes are dejected, suffused with tears, sensitive to the light, red and swollen.

After three or four days we observe an incipient redness of the skin; the fever has acquired much intensity; the cough is nearly incessant; is excited by an annoying tickling in the larynx and trachea, often with considerable hoarseness; the eyes are red, swollen, disposed to water, sensitive to light; there is frequent somnolency and agitation during sleep; the eruption becomes more and more defined, is characterized by its semi-lunar form, and leaves between the red patches certain portions of the skin preserving their natural appearance. In passing the hand

over the skin, it is observed to be slightly elevated, and presents a marbled appearance from the confluence of semi-circular lines. The order in which the eruption is usually observed is as follows: it first displays itself on the face, then behind the ears, on the neck, chest, superior extremities, the trunk, abdomen, and afterwards on the lower limbs. The redness disappears beneath the pressure of the finger, but very quickly returns.

The fever is remittent, with exacerbation in the evening. As soon as the eruption is perfected, the fever gradually subsides, and at this period of the disease the patient frequently exhales an acid odour. Desquamation of the skin is the next phenomenon, but from the delicacy of the scales, it is scarcely observable. The fever has at this time completely ceased, and the cough, which has become moist, gradually declines.

Such is the usual progress of measles, when not in complication with formidable symptoms. The duration of the disease when left to itself, without including the period of convalescence, and when mild, is usually from twelve to fourteen days.

Treated homœopathically, all the stages of the disease are equally defined; but they are, in most

instances, of briefer continuance, and the infection is invariably of milder type.

Pulsatilla is the medicine appropriate to the first stage of the disease: this medicine curtails, renders less painful, and singularly favours the eruption.

Aconite should be exhibited when the disease is ushered in with intense fever, and even during its progress, where the fever is high and where the symptoms indicate an inflammatory condition of the organs of the economy.

Belladonna is applicable in cases where the patient complains of continual thirst; where the tonsils and *velum palati* are inflamed; where the cough is spasmodic, dry, and powerfully affects the chest; the conjunctiva deeply injected; the eyes appear vitreous; are much disposed to tears; also great restlessness of mind; irritability of the nervous system; vigilance; delirium.

Bryonia is very efficient in cases where the eruption has retroceded: it restores it to the surface, and removes the danger attendant upon such internal diversion.

This remedy is more particularly indicated in cases of inflammatory affections of the respiratory organs; or when the eyes are excessively sensitive to the stimulus of light.

Pulsatilla is very useful in cases of diarrhœa and glairy vomitings, by reason of a suppression of the eruption.

Mercury, in certain cases of inflammation of the neck, may be serviceable. The characteristic symptoms which indicate its prescription are swelling of the tonsils, with ulceration; where all the parts are covered with viscid mucus; the gums in a state of inflammation, with salivary secretion more or less abundant.

Mercury, *Pulsatilla*, and *China* are frequently available in the diarrhœas which are consequent upon this disease.

Chamomilla, *Ignatia*, and *Nux* are serviceable in opposing the dry, harsh cough, which continues after the cessation of the disease.

Belladonna, *Cina*, *Hyoscyamus*, and *Conium* are particularly indicated where the cough is spasmodic in its character, and resembles a paroxysm of the whooping-cough.

Practitioners who may have to treat this affection should carefully study the following remedies:

Morbillin, Phosphorus, Nux Vomica, Stramonium, Sulphur, Carbo Vegetabilis, &c.

In an elementary work, similar to the present, we cannot discuss the treatment of the more serious forms of the disease, such as are compli-

cated with nervous fever and cerebral inflammation: such cases must rest in the hands of a practitioner competently versed in homœopathic science. The treatment we have detailed will suffice for ordinary attacks.

The mode of administering the various remedies which have just passed in review must vary according to circumstances. In some individuals, one or two doses will suffice for each of the stages. In others, it will not only be desirable to repeat the dose, but it will be essential to do so. We must dissolve from one to three globules in twelve spoonfuls of water, and exhibit a spoonful every hour, or even every half hour, in cases where the remedies quickly expend their action.

The temperature of the chamber should be moderate.

The beverage should consist of pure water, water-gruel, milk and water, gum-water sweetened with sugar, toast-water, &c.

The nutriment should be very light in the earlier stages of the disease: when the fever runs high, the failure of appetite is evidently an indication of the necessity for still greater abstinence. Soon, however, the appetite will develop itself, and we may gradually revert to a more substantial diet.

The animal powers will be speedily re-established, since the organism has suffered no drawback from the various evacuations prescribed in the allopathic school of medicine.

Scarlatina.

Scarlatina is an epidemic disease, principally incidental to childhood; but all ages are liable to its attacks. Isolated cases of scarlatina have rarely come under observance, and it usually affects the same individual only once in his lifetime.

The febrile symptoms do not set in several days previously to the attack, as in measles: the patient may be apparently in good health to-day, and on the morrow be the victim of disease.

The skin is of a bright scarlet red, similar in colour to a boiled lobster, and the pressure of the finger leaves a white imprint, which instantly afterwards becomes red. This hue of the skin is gradually lost in that of the healthy portion, and is not defined in its margin as in measles. The redness is of frequent variation, sometimes appearing more vivid, at other times paler, in order to reassume a greater intensity.

The skin in true scarlatina is smooth and

glossy; a characteristic which distinguishes it from every other eruption.

The redness commences upon those parts of the body which are uncovered, or but lightly clothed; first the face, throat, breast, hands, feet, &c.; and the redness occupies from three to five days in attaining its full development. When the eruption assumes a different course, it is usually much more serious. Those parts which have been first attacked are invariably the earliest to become pale; and the fever, which is proportionate to the force of the eruption, subsides, and disappears when desquamation supervenes. During the progress of the disease, the skin has remained dry over the parts affected, and it is only at its close that the perspiration is re-established. The desquamation is more decided than in measles, and the epidermis is removed in scales, at times of considerable magnitude.

A characteristic symptom of scarlatina is an affection of the throat. We have observed, that measles is invariably accompanied by cough, and an affection of the eyes; while scarlatina is ever associated with sore-throat, characterized by an intense redness very similar to that of the skin, in which the mucous membrane of the pharynx, mouth, and tongue, more or less participates.

Belladonna is the exclusive remedy for the disease when pure or idiopathic in its character, and free from complication. From one to three globules of belladonna should be dissolved in six spoonfuls of pure water, and a spoonful of the mixture be administered every hour, being guided by the results. We may repeat or intermit the doses at will, and in this manner obviate any loss of time, or, on the other hand, any serious aggravation of symptoms.

Opium is indicated where the disease is complicated with burning heat, a stunning sensation in the head, with somnolency, agonizing disquietude, tossing about from place to place, with vomitings, diarrhœa or costiveness, convulsions, &c. A few attenuated doses of opium will divert the disease into a more kindly and favorable channel.

Ipecacuanha will be of service when the fever exacerbates towards evening; where there is absence of sleep; failure of appetite; and insupportable melancholy, with disposition to tears.

Mercury will be of essential service where the angina accompanying scarlatina is attended with considerable swelling; where the throat appears clogged with viscid mucus, and the parts are really or appear to be ulcerated. In such cases we must not

implicitly rely upon the action of *Belladonna*, but, without loss of time, have recourse to *Mercury*.

Arsenicum is a triumphant remedy in certain cases of angina, especially where the fever has assumed a malignant type. In such instances the throat and mouth are interspersed with numerous ulcerations, exhaling a putrid odour, the animal power is prostrate, the mouth very dry, and the thirst insatiable.

Nux Vomica may succeed to the exhibition of arsenic, where the symptoms have not been entirely removed by the latter agent.

Aconite is frequently of value where the disease is attended either with violent febrile excitement, or high inflammatory action.

Practitioners must consult the following remedial agents where the affection is complicated in its character, viz.

Phosphorus, Rhus, Sulphur, Phosphoric Acid, Helleborus, Cahinca, Coffea.

We must in nowise remit our hygeinic attentions to the patient, and the reader is referred to what has been said upon this subject generally, and to what has been more especially observed under the head of measles.

Small-Pox.

Small-Pox (*Variola*) is an eruptive and contagious disease attacking the patient, in most instances, but once in his lifetime. The affection is regular in its progress, and the stages, of which four are distinguishable, are well defined.

The first is the febrile stage, in which are perceptible upon the skin numerous red spots, proclaiming the disease. The fever is attended with headach, melancholy, ill humour, lassitude, drowsiness, congestion in the head, bleeding at the nose, sleepiness, twitching in the limbs, pendiculation, swelling of the ganglions, and vomiting. The patient complains too of excessive pain in the lumbar region, and this symptom is at times entirely characteristic.

The second is the eruptive stage. The spots or *stigmata* attain a certain height, and present small red elevations, which, after about forty-eight hours, are converted into pustules. They display themselves, in the first instance, on the face; the following day upon the hands, and surface of the body; and on the third day upon the soles of the feet. By this time the pustules are, in most instances, completely formed upon the face, and the patient complains merely of itching

and heat of skin, with pain occasionally in the eyes and in the throat.

The third stage commences after the perfect establishment of the eruption; that is to say, from the fifth to the ninth day. The pustules acquire considerable development, are circumscribed by a red lineation, and are charged with lymph presenting a dark point in the centre, slightly depressed. This lymph, in the first instance limpid, becomes yellow, acquires consistency, is purulent, and the pustules are confluent. When the pustules are numerous, there is considerable swelling, especially about the head and eyes, and the aspect is frequently appalling. At the same time the mucous membrane of the nose, mouth, and pharynx becomes inflamed, and in many instances we find profuse salivation.

Where the eruption is very great, a secondary fever is established: this has been termed the suppurative fever, and, according to circumstances, is more or less violent. The urine becomes thick and clouded, and attended, as it were, with a purulent deposit; while the odour, which is exhaled from the patient, is scarcely supportable.

The fourth stage, that of desiccation and desquamation, terminates the disease. Desiccation commences in those pustules which were first

established, and as soon as this is somewhat confirmed, the suppurative fever decreases, and shortly after gradually subsides. At the same time are observed tumefaction of the face, putrescent odour, and, subsequently, an itching on the skin. The suppuration dries up, and there are formed brown, hard, and dry crusts, which fall off more or less slowly, leaving upon the skin cicatrices, more or less strongly defined. The skin remains for a long time keenly sensible to atmospheric impressions.

This disease is often very formidable. Hitherto we have merely traced the characterizing symptoms; but, superadded to these, we constantly remark an affection of organs mainly essential to life, such as the lungs, brain, stomach, bowels, &c.; and it may be observed, that the more severe the disease, the greater the variety of complication.

The earliest stage of the disease requires the exhibition of *Aconite*, in order to moderate fever and visceral congestion.

Belladonna is indicated where the brain is affected with excessive irritability of the nervous system, and where the eyes are particularly sensitive to the stimulus of light.

Opium is available in cases attended with invincible disposition to sleep.

In the second stage, we shall find *Stramonium* very efficient in facilitating the removal of the eruption, and in limiting its duration. When this stage of the disorder is complicated with gastric phenomena, we may exhibit with advantage *Chamomilla*, *Nux Vomica*, *Crude Antimony*, and *Bryonia*, or even other remedies, the choice of which must be regulated by the characteristic symptoms, and by the changes observable in the moral tendencies of the patient.

Aconite may be productive of much benefit at this period, should fever or inflammatory action be apprehended.

Belladonna is indispensable where the cerebro-spinal economy is affected; where we encounter delirium, ardent thirst, vigilance, and constipation; and where there is burning heat of the body, with coldness of the extremities.

Bryonia and *Aconite* are indicated where the respiratory organs appear to be threatened with inflammation. The eruption, during this period, has frequent tendency to assume a formidable aspect, and either to become identified with putrid fever, or rapidly to disappear. *Arsenic* will prove invaluable in arresting the danger, and in restoring the vital energies which have become nearly extinct.

The third stage may present all the com-

plications of the preceding, and requires, in consequence, a similar mode of treatment, varied with reference to circumstances.

When this period is distinguished by complication, and when we are conversant with its characteristics, there is no remedy more appropriate than *Mercury*, which particularly indicates an affection of the throat, eyes, secretion of saliva, &c.

The desiccating stage requires merely hygeinic observance, and especially cleanliness. In the first instance, ablution must be enforced with warm water, gradually decreasing the temperature, until the water which is used be perfectly cold.

Practitioners will derive much benefit by directing their attention to the following remedies, viz.

China, Coffea, Solanum Mammosum, Sulphur, Variolin, Vaccinin.

Megrim

Is a species of cephalalgia affecting one side of the head, and at times only a small portion, which might, as it were, be covered with the top of the finger, and feeling as though a nail were driven into the part. Occasionally it is but slightly

painful, but in the majority of cases the violence is excessive, and its intensity gradually increases.

There is at first experienced in the affected part a slight pressure, with sense of coldness, similar to what would be induced by pouring cold water upon the part; throbbings, after which the pains become lancinating, tearing, staggering, and insufferable. The immediate seat of the complaint, too, becomes extremely sensitive, and incapable of enduring the slightest pressure.

Occasionally the pain affects the eyelid, whence it ascends to the forehead and temples; or it may proceed downwards to the lip or the gums; or it may station itself in the orbit, causing a profuse lachrymal discharge: the nostril on the side affected is generally obstructed. At times, the megrim extends itself with severity to the occiput and nape of the neck, or affects at the same time the eye, the temple, and the ear. We observe, too, a pricking sensation in the fingers, in the forearm, and in the muscles of the neck; great tension, and strong arterial pulsations in the forehead and in the temples. When the mischief has its seat at the vertex, it occasions a sensation as though the head would split, and it terminates at times by pervading the entire head. In certain cases the countenance is flushed, the eye dreads the light, and it becomes reddened, prominent,

or contracted by spasms. During a violent paroxysm, we may observe slight convulsive movements in the neighbourhood of the eye, or in the nape of the neck, with diffused perspirations which yield no relief.

The paroxysms, which continue frequently for several hours, or even throughout half the day, return in the first instance at irregular intervals; after a while they supervene every day, or every three days, at the same hour; or even every month, with or without shivering succeeded by heats.

This affection is heralded by vertigo, serenity of mind, or unusual loquacity, mental depression, failure of appetite, nausea, colic, acid risings, vomiting, costiveness or diarrhœa, with swelling of the lower abdomen, and terminating either in sleep, perspiration, vomiting, or tears.

This disease is very obstinate, and frequently affects women, sometimes from their infancy, but more usually from the period of puberty, to the age of fifty or sixty years. The accidental circumstances which produce, aggravate, or modify other pains in the head, possess no influence whatever over megrim.

The almost invariable cause of this malady is a psoric tendency in the constitution, and the patient must necessarily submit to a tedious anti-

psoric treatment in order to accomplish a cure. When this psoric condition has been effectually checked, we may derive very beneficial results from *Nux Vomica*, *Ignatia*, and *Pulsatilla*.

Under certain circumstances, where it is evident there can be no psoric principle, we must commence our treatment by the medicines of which we shall presently speak.

Nux Vomica will stand in the foremost rank, more particularly when the pain commences in the morning on rising, becomes aggravated, and where we find a state of habitual constipation.

Pulsatilla is eligible in hemicrania affecting the left side of the forehead, with throbbing and shooting, night and morning, increased by stooping in the room, or on moving the eyes; also where we find stomach-ach, with bilious, acid vomitings, with griping and pinching pains in the abdomen.

Pulsatilla and *Ignatia* are equally valuable to succeed *Nux Vomica* as an intermediate remedy, since they possess an action very analogous to that of the latter medicine.

When *Megrim* presents itself in an extremely violent form, we must have recourse to *Coffea*, which, should it fail in its complete removal, will at least render the pain more tolerable.

When the megrim proceeds from an abuse of

coffee, *Chamomilla*, *Nux Vomica*, and *Ignatia* will be the most appropriate remedies.

Belladonna is especially suitable where the pain in the head is lateral; extends to the orbit, and to the bones of the nose; where the pain is characterized by pressure and darting, with a sense of boiling, as of a liquid, in the head; where the pain is aggravated by the slightest motion of the head, or of the eyes; and where these organs are extremely sensitive to the slightest ray of light; where every description of noise is insupportable; and where we find strong arterial pulsation; also where the pains are characterized by violent shooting, nearly depriving the patient of his senses.

Colocynthis is appropriate where the pain is compressive, twitching, or pinching; and where the pains are increased either by stooping or lying upon the back.

In an elementary work we must content ourselves with directing attention to cases most frequent in occurrence, and refer the practitioner to the following remedies in the more rare or complicated affections.

Sepia, *Nitric. Acid.*, *Zincum*, *Petroleum*, *China*, *Bryonia*, *Veratrum*, *Capsicum*, *Ipecacuanha*, *Cicuta*, *Aurum*, *Arsenic*, *Mangan. Acetat.*, *Causticum*, *Rhus*, *Guaiacum*, *Arnica*, *Hyoscyamus*, *Lycopodium*, *Phosphorus*, *Conium*.

Odontalgia, or Tooth-ach.

This affection in most instances originates in a cold, or access of blood to the head; and among hysterical women, in excitement of the nervous sensibility. In cases which are very obstinate and long existing, it is the result of a psoric tendency, and a treatment, to be crowned with success, must be directed to the remote cause in which it takes its rise.

Homœopathy is in possession of a variety of efficient means which are applicable to the different peculiarities exhibited by affections of the teeth.

Odontalgia, resulting from a cold, affecting merely a hollow tooth; drawing and boring in its character, and accompanied by occasional and isolated shooting pain, in which the entire economy participates; and where we meet with at times a painful swelling of the gums, will yield to the exhibition of *Nux Vomica*. This result will more particularly take place where the pain supervenes early in the morning while in bed; where mastication is prevented; and where aggravation is induced by contact with cold air. This remedy is particularly suitable to persons of

lively temperament, and accustomed to the use of coffee and spirituous liquors.

Mercury is applicable where the drawing or boring pain results from a cold; where we find swelling of the gums and of the cheek; shaking of the teeth; where the pains increase after a repast, or after having drunk anything hot or cold; where the salivary glands are affected, and secrete an excess of saliva, causing the patient to spit frequently; and where there is ulceration of the gums.

Hyoscyamus is the most effective remedy against odontalgia exhibiting itself in the morning, especially after a cold, or under the influence of cool air, with considerable flow of blood to the interior of the head; throbbing and plucking pain from the cheek to the forehead; and thrilling in the tooth, which is loosened; tearing pain in the gums.

Aconite will be found very useful in throbbing odontalgia, originating in a cold, and accompanied by determination of blood to the head, and burning flushes of the face.

The north-pole of the magnet is appropriate in rheumatic odontalgia, where all the carious teeth are affected, and where we observe the following phenomena:

The gums swollen, sensitive, and painful to the

touch ; isolated paroxysms ; spasmodic pressure ; tearing, searching pains ; burning lancinations ; pain in the jaws and the incisor teeth, induced by respiration.

The magnetic pole should be touched during one or two minutes only, and the process repeated one or more times according to the result.

Rhus effects the removal of tearing odontalgia, of which the application of warmth will occasion a decrease ; it is eligible, too, where the pain resembles that of a wound ; where the shootings are spasmodic ; and where we find twitchings and tearings, aggravated towards night, or in the open air, and alleviated by warmth.

Staphysagria is a suitable remedy where the odontalgia, ceasing to affect the carious tooth, communicates itself to the adjacent ones, or to those on the opposite side, and extends even to the ear, accompanied by pressive pain in the temple on the same side. Also, when the suffering is experienced on the air entering the mouth ; gnawing pain in the inferior incisors, especially in the night ; twitching pain in the teeth after having eaten, and aggravated in the open air.

Belladonna is suitable in odontalgia after a cold, especially among women ; and when the pain is rendered insufferable from the touch, the air, and the action of eating. This pain is of a

drawing character, and extends from the ear to the hollow teeth of the upper jaw, where at times it becomes thrilling; is aggravated in the night; when the pain is not increased during a meal, but becomes increased shortly after it.

Bryonia is available in cases where the pain impels the patient to go to bed, or where it is exacerbated by heat at night; on lying down on the side not affected, or relieved by lying on the side which suffers; spasmodic twitchings, with a feeling as though the teeth were too long, and, as it were, loosened, more particularly during and after a repast; boring and plucking pain, decreasing in the open air, and by cold water in the mouth; drawing pain, at the time, in the cheek, and pinchings in the ear, continuing from night till morning.

Chamomilla may be prescribed in odontalgic affections resulting from a cold, with plucking pains in the jaws; after drinking and eating; especially after hot things, coffee, &c.; pains in the teeth extremely violent in the night, with swelling of the cheek, aggravated by cold drinks, and rendered intolerable by the heat of the bed, swelling of the gums, &c.

These pains in the teeth presenting a vastly numerous variety, we have confined ourselves to those of more frequent occurrence, referring the

practitioner to the *Materia Medica* for those cases which are not comprised among the remedies we have pointed out.

Otalgia, or Ear-ach.

We thus denominate a violent pain affecting one of the ears, or both of them at the same time, and which frequently embraces the adjacent parts, where it is accompanied by pains in the teeth, or other rheumatic affections, and of which the most usual cause is a cold. We rarely observe either redness or external swelling; the patient complains of darting, throbbing, quivering pain, with tinkling and humming in the ears. He experiences great agitation, anxiety, vigilance, pains in the head, and fever. Frequently, after the lapse of some days, there is established a watery or purulent discharge, which diminishes the suffering.

Mercury should be administered where the pain experienced in the interior of the ear is tearing or shooting, with sensation of cold or of squeezing, and affecting also the cheek; where there is perceptible inflammation in the auditory canal, and discharge of pus, with ulceration of the conduit.

Arnica is eligible where there is general sensi-

bility of the nervous system, upon the slightest exciting cause; pressure; dartings in and behind the ears; tearing pains, with sensation of heat; sensibility to sounds which are loud and sonorous.

Chamomilla is appropriate where the ear-ach proceeds from cold, or from an arrest of perspiration; where there is tearing pain in the ears, and as far as the lips; lancinations strong and isolated; with uneasiness and dryness of the ear.

Pulsatilla is suitable to persons subject to rheumatic affections, and who suffer from pains in the ears, which are usually characterized by isolated spasms, with plucking sensations externally and internally; heat, with external redness of the entire ear; painful sensation in the ear, as though something would be discharged from it; want of sleep; and, among sensitive persons, an inclination to tears, cold shiverings, &c.

Belladonna is applicable to otalgia with perforating and shooting pain in and behind the ear; plucking and shooting in the parotid glands, even to the pharynx; when the suffering is more lively within than without; and especially where the patient experiences cerebral and sympathetic symptoms, such as delirium, convulsions, syncope, aphonia, &c.

Palpitations.

Palpitations, in a majority of instances, are merely an accessory symptom of maladies more or less important. Such cases demand, in consequence, the treatment of the affections of which they are merely symptomatic.

In certain individuals of full and plethoric habit, or where other causes may induce an afflux of blood to the chest, we sometimes remark palpitations without any other affection.

To remedy this ailment we must carefully avoid stimulating and spirituous drinks, and take a few doses of *Aconite*, at some days' interval. This medicine quiets the circulation in a very powerful manner.

After the *Aconite*, we may administer one or more doses of *Nux Vomica*, provided the mischief be not entirely removed.

Chamomilla is a suitable remedy where the palpitations supervene upon a fit of passion, or violent anger, producing great distress.

Nux Vomica is especially indicated where the exciting causes of the palpitations are stimulating drinks, coffee, &c.

Pulsatilla is a remedy suitable to women of very sensitive and delicate temperament, and characterized by sweetness of disposition.

Whitlow (Paronchia)

Is a local inflammation, having its seat in most instances at the extremity of one of the fingers. As its symptoms, we find swelling, violent pain, afterwards suppuration; it frequently reappears among persons who have once been affected. It sometimes displays itself in the absence of all external cause; sometimes after a cold, a contusion, burn, or any other lesion; after the removal of the epidermis at the roots of the nails, from cutting the nails too near the quick, or from having touched any corrosive liquid.

The pain is moderate when experienced in the cellular tissue; more vivid when affecting the periosteum; and intolerable when the inflammation affects the immediate envelope of the tendons.

It is not unusual to find this malady producing, among persons of irritable temperament, either convulsions or fainting fits.

Whitlows yield readily to *Mercury*; but if such means should not suffice, we may subsequently have recourse to *Sulphur*. The *North Pole of the Magnet* is equally powerful in these affections.

Hepar Sulphuris and *Rhus* are also valuable remedies in these cases; but when they return

frequently after the application of the *Magnet*, and the exhibition of *Mercury*, it will be important to administer *Sulphur* and *Silicea*.

Chicken-pox.

This eruptive disease presents no danger in itself, and frequently attacks children, in districts infected with small-pox. In its external aspect, it frequently bears much resemblance to the last-mentioned disease. It may be distinguished, however, by the more pointed form of the pustules; the irregularity of their appearance; the absence of smell peculiar to variola; by the mild character and rapid progress of the disease; and by its briefness of duration. It more especially affects the face; is accompanied by slight cold in the head, trifling cough, and slight febrile excitement; and terminates generally in about seven days.

We administer *Aconite* at the moment of fever; and, if there be any agitation or anxiety, we have recourse to *Coffea*. Should spasmodic symptoms supervene among children, and especially during dentition, we must have recourse to *Chamomilla*.

Belladonna must be prescribed if the child raise the head with difficulty; if the head and hands

are hot; and if there be agitation, and griping pains.

Aconite, at the commencement of the evil, will curtail its duration.

Mercury is applicable where the pustules secrete matter. This remedy, however, is more especially applicable in variola and chicken-pox at the period of suppuration.

Rheumatism.

This affection has its seat in the membranous structure; in the tendons, ligaments, periosteum, and even in the bones; and chiefly in the muscles and their envelope.

Rheumatism frequently confines itself to a single muscle, tendon, or membrane, the other muscles retaining their functions intact. In such cases, certain movements and certain positions of the body procure solace or tranquillity to the patient.

Where the rheumatism affects a large surface, or an assemblage of external muscles, or articulations, it is impossible to move without experiencing the most excruciating pain.

The pain is in most instances tearing, darting, drawing, and burning; sometimes cutting, exten-

sive or pressive; constrictive, or consisting in a sense of dulness, and seldom assuming a pulsative character.

The suffering is at times extremely violent, aggravated by the slightest external cause, or by the most trivial movement of the affected muscle.

It is alleviated by rest, leaving merely a painful sense of weakness, but it is scarcely ever entirely suppressed; and even under the most favorable circumstances, the patient experiences occasional shooting pains. The suffering is not confined to a single locality, but is diffused throughout the whole affected region. The heat of the bed increases acute rheumatism; the pain subsides, quickly changes its situation; returns afresh; and in this manner is limited neither in its seat nor in its duration. It rarely affects the internal organs, and evinces a great tendency to occupy its original locality.

Among the articulations in which rheumatism produces an effect analogous to that of luxation, we may instance the shoulder, the knee, and the arms; it less frequently attacks the wrist, and still more rarely the joints of the fingers and great toes. The progress of the disease is slow, and generally presents no serious danger.

At times it is dispelled very readily; but there remains a predisposition to its return under the

influence of cold, or of any other adventitious cause.

It affects persons even of the most robust habit; of both sexes; and of every age and condition. In youthful persons it more particularly affects the superior parts of the body, from the head to the hands; and among persons more advanced in years, it is felt in the back, the loins, the haunches, and the thighs.

Rheumatism is in most instances preceded by premonitory symptoms; it is frequently sudden in its access, and principally results from cold, or from a transition from a higher to a lower temperature: weakness and sensibility of the skin indicate a predisposition: and it may occasionally result from some interference with perspiration.

Chamomilla is a suitable means of relief, when the rheumatism presents the following symptoms:

General uneasiness; heaviness in the limbs; tearing pains in different parts of the joints, aggravated at night, and diminishing on sitting up in bed; all the limbs feel as though paralyzed; pain in the loins; tooth-ach, affecting the entire side of the face, and extending to the ear.

Pulsatilla is effective where the rheumatism is characterized by the following phenomena:

Drawing pains, exacerbated towards evening, planting themselves in the limbs, and particularly

in the muscles which border upon the articulations; pain in the joints of the feet; sense of general debility; disturbed sleep; shiverings; pressive pain in the lower part of the abdomen during motion; cough; sneezing; pains frequently changing their position.

Bryonia is applicable under the following circumstances:

Aching pain in the joints of the arm, the back, nape of the neck, and in the breast; and experienced chiefly during motion or inspiration, perceptible even when seated or lying down, and aggravated by moving or walking about; pains in the loins; vivid, drawing, tearing pain in the tendons of the muscles, and at the point of union with the extremity of the bones, especially after rising in the morning; pain in the limbs during motion, as though they were dislocated; stiffness of the body when moving, especially after mid-day.

Nux Vomica is applicable to the following train of symptoms:

Tension and pressure in the external parts of the chest; shooting pains in the muscles of the chest, and between the shoulder-blades, during movement and respiration; pain resembling a cracking in the shoulder-joint and in the shoulder-blade; tearing pain at the nape of the neck; pain

in the cervical vertebræ; stiffness of the back; violent pains in the loins; agitation; shiverings; coldness of the feet and hands; disposition morose and querulous.

Aconite is indispensable where the rheumatism presents itself under an inflammatory type; and where the parts are red and swollen. When these symptoms have been checked by *Aconite*, and where the fever has been much subdued, we may administer *Antimonium Crudum*, which may be repeated as circumstances shall indicate. After this remedy, or even after the *aconite*, we may exhibit *Bryonia*, particularly should the pain reappear, or be aggravated by any motion of the affected part.

Rhus Toxicodendron is of signal service where the rheumatism is characterized by the following indications:

Tension, drawing, and tearing in the limbs; more violent during rest, with sensation of torpor and dulness in the affected parts after motion; pains of bruising, or, as though the flesh were torn from the bones; pressive, drawing pain in the periosteum, as though the bone were being scraped; where patients suffer more particularly at night, and are relieved by motion.

Dulcamara is of much value in rheumatism arising from a cold, and particularly from a cold

dampness in the atmosphere; when the pains are characterized by shooting, drawing, and plucking, and have their seat in the limbs; when the upper extremities are, as it were, paralyzed or cold; feel as though they had received a severe contusion; and are more especially experienced in the night and during rest.

The practitioner may consult the following medicines with considerable advantage:

Belladonna, Ignatia, Phosphorus, Sulphur, Thuja, Veratrum, Arnica, Calcarea, Mercurius, Cocculus, Silicea, China, Arsenic, Rhododendron, Tinctura Acris.

Cold in the Head

Displays itself after a cold by a particular sensation in the nasal fossæ, commencing with dryness and swelling in the part, a little increased during inspiration; frequent sneezing, nasal intonation, and loss of smell. The discharge is frequently watery, saltish, falls drop by drop, and produces excoriation, dryness, and crusts upon the nostrils and lips. The mucous secretion comes from the frontal cavities, the upper jaw, and from the nose, where the patients experience a sensation more or less painful. This affection

is occasionally preceded and accompanied by dryness, and by a disagreeable smell of dust, with violent pressive pain at the root of the nose, and in the frontal sinus, constituting that state of things which has been named a cold in the head.

Cold in the head consists essentially in an irritation of the mucous membrane of the nasal fossæ. When this irritation is very violent, it is accompanied by fever.

Nux Vomica is the appropriate remedy when we find obstruction, with great dryness of the mouth; heat of face increased towards evening; itching in the nose; painful sensibility of the interior of the nose; obstruction during night only; secretion from the nose during the day; heat and confusion in the head; general languor, grief, and vexation.

Pulsatilla exercises a powerful control over the following symptoms:

Tickling sensation in the nose, as from very fine snuff; very violent sneezing; loss of smell; discharge of blood from the nose on using the handkerchief; secretion of fetid mucus from the nose; painful ulcerations in the nostrils; sensibility of the eyes to light; pressive head-ach; broken sleep; melancholy; inclination to weep; shiverings; hoarseness.

Chamomilla is applicable to cold in the head, presenting, as its characteristics, a discharge of

mucus from the nose; the nostrils ulcerated, chapped, and inflamed; pain in the lips, which desquamate; somnolency; confusion of the head; giddiness; one cheek red, the other pallid; shiverings; and intense thirst. This medicine is particularly appropriate to children.

Mercury renders effective service when there is excessive discharge; the nose excoriated both within and without; and where we find rheumatic pains in the limbs.

Under peculiar circumstances the preceding medicines may fail, the practitioner will therefore refer to the following:

Arsenic, Bryonia, Belladonna, Aurum, Meze-reon, Ammonium, Lycopodium, Sulphur, Natrum, Antimonium Tartarizatum.

Bleeding from the Nose (Epistaxis).

Persons who have frequent determination to the head, children, and young people are particularly liable to this inconvenience. The blood is usually discharged from one nostril only; sometimes during sleep, it flows into the pharynx, and causes cough and sickness, so that the disease may be mistaken for vomiting or spitting of blood. If the bleeding be very trivial, we may leave nature

to her own resources; since her own unaided efforts will remedy the evil: but when the blood is discharged in considerable quantity, and when the hemorrhage is frequently renewed, it is necessary to interpose a remedy, lest it should become prejudicial to health.

Arnica is especially suitable when the flow of blood is preceded by prickings in the nose and forehead, as from some insect; when the nose is hot, the blood quite fluid, and of a bright red colour.

Pulsatilla is appropriate for adult females, when the monthly periods last a shorter time than is consistent with health, and are less abundant; and particularly when the moral temperament is sweet and placid.

Crocus is suitable to women who are irritable and sensitive; whose blood is in a state of ebullition; who feel a diffused heat, and discharge thick and dark-coloured blood from the nose.

Mercury is applicable to epistaxis when the blood coagulates in its flow, and remains suspended. This remedy is similarly appropriate for children, and for parties disposed to inflammatory affections.

China may be administered subsequently to the discharge, in order to prevent its return; we may commence its exhibition twenty-four hours after

the hemorrhage; and, according to the result, may repeat the same medicine two or three times.

Belladonna is very useful where we meet with discharge of blood from the nose, with determination to the head, evincing itself by a swelling of the veins; darting pain, returning in paroxysms, in one side of the head, and aggravated by noise, light, or motion; scintillations before the eyes; dimness of vision; tingling in the ears; syncope; tendency to sleep.

Under certain circumstances the following medicines will be similarly appropriate:

Aconite, Arsenic, Ignatia, Nux Vomica, Chamomilla, Cina, Carbo Vegetabilis, Bryonia, Moschus, Rhus, Secale Cornutum, Sulphur, Sepia, Graphites, Lycopodium.

Stomacace (Ulcerative Inflammation of the Mouth).

We thus denominate an affection which commences with an attack on the front part of the mouth, and afterwards extends to the uvula and palate. The earlier symptoms are stiffness and twitching in the throat; a taste of copper, with burning; heat, redness, swelling, and extreme sensitiveness of the gums, interior of the lips, cheeks, tongue, and palate. To these phenomena,

are subsequently united a cadaverous and nearly intolerable stench in the mouth; painful enlargement of the glands of the neck; profuse discharge of saliva or viscid and fetid mucus from the mouth. The gums are spongy, extremely swollen or retracted; the teeth painful, foul, and oftentimes loose; mastication, swallowing, speech, and respiration, attended with much suffering. At length upon the affected parts are formed small flat ulcerations, very sensitive and obstinate. These ulcerations extend themselves; the inferior parts are foul, resemble lard, and are fungous; the edges are soft and slightly elevated, inflamed, and uneven. In many cases we observe fungoid excrescences, differing in form, colour, consistency, and size. Very frequently there flows from the parts which are swollen and ulcerated, particularly the gums, and especially when touched, a putrid sanies or blood of a deep-red colour. At the same time there is much debility, and very frequently much febrile excitement.

Mercury is a very efficient weapon with which to oppose this malady: a few doses will in most instances suffice for its ready removal.

Nux Vomica, in certain cases, is very useful; in derangements, for instance, in the digestive functions; constipation, &c.; and where the affection is characterized rather by an inflam-

mation and swelling of the membranous parts than by an ulcerative inflammation with discharge of blood or sanious matter.

Cough.

Catarrhal Cough, of which the origin is in most instances a cold, will yield very frequently to the application of *Nux Vomica*, more particularly when the cough is excited by a sense of roughness and scraping in the throat, with titillation in the palate; if the cough be constant, exhausting, and cause pain in the head, which feels, at each access, as though it would split; if, as is still less frequent, a violent pain, as of a bruise, be experienced in the lower part of the abdomen; beneath the ribs, after coughing, a pain as from a blow and if respiration be impeded during the night. Also, when in addition to these symptoms we find the sensation of a heated substance, pressing upon the chest; when the cough is more especially troublesome in the morning towards four or five o'clock; when it is excited by speaking, motion, &c.

Chamomilla is particularly adapted to children when the cough results from a cold; when it is strong and dry even during sleep; when it is excited in the day-time by a continual titillation in the trachea behind the hollow of the throat;

when it seems, in coughing, as though something were rising in the throat, and likely to take away the breath; when the cough is excited by a fit of passion, especially in children.

Hyoscyamus may be employed with advantage when the cough is dry, frequent, and especially during the night, so as to prevent sleep; if it be almost constant when lying down, and cease when sitting up in the bed; when the patient is compelled to resume the latter position at brief intervals, in order to obtain some slight relief; when to these symptoms is united a tickling irritation in the trachea, with a species of convulsive cough.

Belladonna is often very valuable in dry spasmodic cough, with asthmatic breathing, occurring at night after a cold; pain beneath the sternum at night, or after a meal, with difficulty and failure of respiration; pain in the hypochondria; constriction in the chest; swelling of the abdomen; pain in the loins, with disposition to be alarmed; cough, with stertorous breathing and listlessness.

Aconite is serviceable in cough, at the termination of which the chest feels lacerated; where the fits are lasting; where the paroxysms are short, and leave behind them a sensation of burning in the chest. *Aconite* is more par-

ticularly indicated where the disease more nearly approaches the inflammatory state.

Ignatia is especially applicable to a dry cough, proceeding from the trachea, with pain in the lower region of the abdomen, and tightness across the chest; and where the cough exists both day and night with similar intensity.

Ipecacuanha should be administered in dry, spasmodic, shaking cough, with suspension of the breath, and loss of sense in dry cough, with painful jerkings in the head and stomach; nausea and vomiting; diffused perspiration; constant obstruction of the nose, with loss of smell; mucous expectoration unpleasant to the taste; tickling in the throat, which is, as it were, constricted; sensation as of a wound in the chest, and aggravated by cold air.

Pulsatilla should be prescribed where the cough is distressing both by day and night, with pain affecting both sides, and expectoration after a long-continued fit of coughing; profuse nocturnal sweats; thirst; small and frequent pulse; emaciation; loss of appetite; cough in the morning, with tremor of the whole body; water discharging from the mouth, with inclination to vomit; expectoration of mucus, yellow, and afterwards salt or bitter to the taste, and producing nausea, sputa occasionally tinged with blood.

In certain cases the following medicines may be successfully administered:

Phosphoric Acid, Aconite, Arsenic, Bryonia, Drosera, Mercurius, Sepia, Spongia, Stannum, China, Sulphur, Conium, Lactuca Viridis, Arnica, Calcareo, Lycopodium.

The study of the *Materia Medica* will explain under what circumstances either of the above remedies will be appropriate.

Swelling of the Parotid and Submaxillary Glands.

This malady occasionally precedes affections of the teeth, with which it will disappear. More frequently it has its immediate origin in a cold, or in atmospheric changes. It may attain considerable importance, attack both sides of the face, and degenerate, at times, into a hard suppurative tumour. When the disease is violent it is accompanied by an erysipelatous inflammation, together with febrile excitement.

Mercury is the most appropriate remedy, even at all periods of the disease, as also when the tumour enters upon its suppurative stage. The remedy may be repeated several times if desirable, at some days' interval, or every day, in water.

Belladonna may be administered where we find an erysipelatous inflammation; when the pain subsides we may have recourse to *Hepar Sulphuris*, which will perfect the cure.

Dulcamara will render powerful aid against an induration of the glands, when the disease has not been checked by the exhibition of *Mercury* and *Hepar Sulphuris*.

Worms.

Worms are engendered under the influence of uncleanness, an unwholesome character of the food, unhealthy atmosphere, moisture, and by every circumstance which may favour the accumulation of mucus in the intestinal canal.

The species of worm most frequently encountered, are the *Ascarides*, the *Lumbricales*, and the Solitary or Tape-worm. Women and children are usually more subject to this visitation than men.

The *ascarides* bear a resemblance to the mites in cheese; some of them, however, attain the length of upwards of an inch. These worms are frequently met with in children, and excite a very unpleasant itching in the alimentary canal, with

frequent and pressing inclination to relieve the bowels, together with other symptoms.

The lumbricales resemble the earth-worm. They are white, and have, at the top of the head, a circle studded with slight eminences or warts. These worms attach themselves in clusters to different portions of the intestines, beneath the parietes of which they create, in moving about and sucking, and especially after having eaten sweet food, an irritation which gives rise to painful griping, more particularly in the umbilical region. The ordinary indications of their presence are the following :

Nausea ; accumulation of watery saliva in the mouth ; fetid breath ; face pale and swollen ; a blue or brownish hue beneath the eyes ; dilatation of the pupil ; broken sleep ; urine thick and white ; and the lower part of the abdomen hard and tender to the touch.

The solitary worm is flat and whitish, frequently of considerable length, and comprising a vast number of articulations.

It is sufficiently difficult to recognize its presence, since—if it occasion at times distressing symptoms, extraordinary sense of pressure, coldness, undulatory motion in the lower region of the abdomen, lively pain, cramps, epileptic and convulsive fits, &c.—it at other times gives rise to none of these sensations. In this case, the only

positive indication is the discharge of certain detached portions of the worm.

An antipsoric treatment is indispensable, in order effectually to check the disposition which favours the generation of worms. Since the conditions which we have pointed out are, in most instances, the accidental causes giving rise to worms, it is equally true that there exists at the same time an internal cause peculiar to the individual, and this cause is the psora, which must be subdued.

Such course of treatment bears reference to chronic maladies, with which it is not the purport of this treatise to interfere.

As regards the inconveniences to which parties annoyed with worms are liable, we must meet them, according to circumstances, by the following remedies:

Against *Ascarides* we may prescribe Aconite, Nux, Mercury, Ignatia, Ferrum, Valerian.

Against *Lumbricales*, Nux, Cina, Belladonna, Mercury.

Stramonium will relieve the colic induced by the presence of worms.

Cicuta Virosa is very efficient in fevers excited by worms, and accompanied by cramps and violent pains in the abdomen,

Tinctura Sulphuris removes the itching in the

rectum, resulting from a similar cause. *Filix Mas*, administered from time to time, will in most instances relieve the phenomena, frequently painful and distressing, induced by the presence of the solitary or tape worm.

Vomiting.

The treatment of vomiting must be regulated by the exciting cause, since the latter serves perfectly well to decide its character.

If the vomiting proceed from a superabundance of food, it will be unnecessary to offer any opposition, since it is an effort of nature to relieve the stomach from the substances with which it is overcharged. When the sickness does not establish itself voluntarily, we may assist the vital force and promote vomiting by draughts of warm water, coffee, &c.

If, after an abundant vomiting, there remain aversion, nausea, and heaving of the stomach, we may check such symptoms with one or two doses of *Antimonium crudum*.

Pulsatilla may be selected in preference, where the vomiting has been induced by very gross or fat articles of diet.

Chamomilla is appropriate to cases of vomiting,

excited by a violent fit of passion, and especially where we find, as its accessories, a bitter taste in the mouth; bilious and bitter risings; where the ejected matter is green, bilious, and causes a grating sensation in the throat; if the patient complain of fulness and pressive pain in the stomach; general languor; total loss of appetite; anxiety; heat; excessive thirst; vertigo; and semi-lateral pains in the head.

Bryonia is a remedy which is applicable where the fit of passion continues during the vomiting, and where we perceive a sensation of coldness united to the symptoms above detailed, as referrible to chamomilla.

Aconite is the most efficient remedy in vomiting excited by passion, and accompanied by fear.

Valerian is suitable for children who are frequently attacked with vomiting, occasioned by the presence of worms, and with which are ordinarily associated pains in the abdomen, frequent nausea, excess of saliva, with vapid and ropy taste; pallor of complexion, discoloration of the lips, with general coldness, and accession of debility. These symptoms, however, being dispersed by the medicine, it is essential that the children be subjected to an antipsoric course of treatment, which alone will have the power of confirming the cure.

Pulsatilla is very serviceable in vomitings of viscid whitish mucus, to which many persons of irritable temperament, and subject to spasmodic affections, are disposed. These vomitings are usually preceded by vertigo, pains in the abdomen, by faintings, and in most instances are followed by diarrhœa.

Should *Pulsatilla* fail in its effect, we may administer *Cocculus*. The latter remedy is especially appropriate where the vomitings have resulted from carriage exercise.

The following medicines are frequently of great value in vomitings; and the practitioner cannot too sedulously study their properties in the *Materia Medica*.

Arsenic, Digitalis, Ipecacuanha, Nux Vomica, Veratrum, Silicea, Graphites, Phosphorus, Lycopodium, Sulphur.

Printed by C. Adlard, Bartholomew Close.
